

**SOME HIGHLY  
COMPUTATIONAL  
PROBLEMS CONCERNING  
INTEGER POLYNOMIALS  
OF SMALL NORM.**

PETER BORWEIN

Simon Fraser University Centre for  
Constructive and Experimental  
Mathematics

<http://www.cecm.sfu.ca/~pborwein>

Typeset by  $\mathcal{AM}\mathcal{S}$ -T<sub>E</sub>X

The PTE problem is solved by two sets of  $n$  integers satisfying any of the following:

$$\sum_{i=1}^n \alpha_i^j = \sum_{i=1}^n \beta_i^j \quad j = 1, \dots, n-1$$

$$\prod_{i=1}^n (x - \alpha_i) - \prod_{i=1}^n (x - \beta_i) = C$$

$$(x-1)^n \mid \sum_{i=1}^n x^{\alpha_i} - \sum_{i=1}^n x^{\beta_i}.$$

The conjecture due to Wright (and others) is that it is always possible.

Almost equivalently (though not quite obviously) find a polynomial with coefficients  $\{0, -1, +1\}$  with

$$\|p\|_{L^2\{|z|=1\}} = \sqrt{2n}.$$

## Partial History.

- Euler
- Prouhet (1851)
- Tarry (1910) - Small Examples
- Escott (1910) - Small Examples
- Wright, Fuchs (1935) - Easier Waring
- Letac (1941) - Size 9 and 10
- Gloden (1946) - Size 9 and 10
- Smyth (Math Comp. 1991) - Size 10

An **even ideal symmetric solution** of size  $n$  is of the form

$$\{\pm\alpha_1, \dots, \pm\alpha_{n/2}\}, \{\pm\beta_1, \dots, \pm\beta_{n/2}\}$$

It satisfies

$$\sum_{i=1}^{k/2} \alpha_i^{2j} = \sum_{i=1}^{k/2} \beta_i^{2j} \quad j = 1, \dots, \frac{k-2}{2}$$

An **odd ideal symmetric solution** of size  $n$  is of the form

$$\{\alpha_1, \dots, \alpha_k\}, \{-\alpha_1, \dots, -\alpha_k\}$$

and satisfies

$$\sum_{i=1}^k \alpha_i^j = 0 \quad j = 1, 3, 5, \dots, k-2$$

Listed below are ideal symmetric solutions for sizes  $2 \leq n \leq 10$ , the odd symmetric solutions are all perfect. These solutions are listed in abbreviated symmetric form. For example the solution for size 6 is

$$\{\pm 4, \pm 9, \pm 13\}, \{\pm 1, \pm 11, \pm 12\}$$

and the solution for size 5 is

$$\{-8, -7, 1, 5, 9\}, \{8, 7, -1, -5, -9\}.$$

$$2 : \{3\}, \{1\}$$

$$3 : \{-2, -1, 3\}$$

$$4 : \{3, 11\}, \{7, 9\}$$

$$5 : \{-8, -7, 1, 5, 9\}$$

$$6 : \{4, 9, 13\}, \{1, 11, 12\}$$

$$7 : \{-51, -33, -24, 7, 13, 38, 50\}$$

$$8 : \{2, 16, 21, 25\}, \{5, 14, 23, 24\}$$

$$9 : \{-98, -82, -58, -34, 13, 16, 69, 75, 99\}$$

$$9 : \{-169, -161, -119, -63, 8, 50, 132, 148, 174\}$$

$$10 : \{436, 11857, 20449, 20667, 23750\},$$

$$\{12, 11881, 20231, 20885, 23738\}$$

$$10 : \{133225698289, 189880696822, 338027122801, \\ 432967471212, 529393533005\},$$

$$\{87647378809, 243086774390, 308520455907, \\ 441746154196, 527907819623\}$$

**Size 5.** The following is a one parameter example of size 5.

$$F_5 :=$$

$$\begin{aligned} & (t + 2m^2) (t - 1) (t + 2m^2 - 1) \\ & (t - 2m^2 + 1 - m) (t - 2m^2 + m + 1) \\ & - (t - 2m^2) (t + 1) (t - 2m^2 + 1) \\ & (t + 2m^2 - 1 + m) (t + 2m^2 - m - 1) \end{aligned}$$

This expands to

$$\begin{aligned} F_5 := & -4m^2(m - 1)(2m + 1)(2m - 1) \\ & (m + 1)(2m^2 - 1) \end{aligned}$$

**Size 6.** It is possible to completely solve the even symmetric problem of size 6 in Maple. Basically one just uses “solve”. it gives the following general solution (translated with  $a_6 = 0.$ )

This gives as rational solution of size 6:

$$\{a_2 = a_2, b_1 = b_1, b_3 = b_3,$$

$$b_2 = \frac{a_2^2 - a_2 b_3 + b_1 b_3 - a_2 b_1}{-b_1 - b_3 + a_2},$$

$$a_1 = 2/3 \frac{a_2^2 - b_3^2 - b_1^2 - b_1 b_3}{-b_1 - b_3 + a_2},$$

$$a_3 = \frac{-b_1^2 - b_1 b_3 + a_2 b_1 + a_2 b_3 - b_3^2}{-b_1 - b_3 + a_2} \}$$

The following is a simple two parameter example of size 6.

$$F_6 :=$$

$$\begin{aligned} & \left( t^2 - (2n + 2m)^2 \right) \left( t^2 - (nm + n + m - 3)^2 \right) \\ & \left( t^2 - (nm - n - m - 3)^2 \right) \\ & - \left( t^2 - (2n - 2m)^2 \right) \left( t^2 - (-nm + n - m - 3)^2 \right) \\ & \left( t^2 - (-nm - n + m - 3)^2 \right) \end{aligned}$$

On expansion one sees that

$$\begin{aligned} F_6 := & -16nm(m-1)(m+3)(m-3)(m+1) \\ & (n-1)(n+3)(n-3)(n+1) \end{aligned}$$

**Size 7.** Gloden simplified.

$$(t - R_1) (t - R_2) (t - R_3) (t - R_4)$$

$$(t - R_5) (t - R_6) (t - R_7)$$

$$- (t + R_1) (t + R_2) (t + R_3) (t + R_4)$$

$$(t + R_5) (t + R_6) (t + R_7)$$

where

$$R_1 := - (-3 j^2 k + k^3 + j^3) (j^2 - k j + k^2)$$

$$R_2 := (j + k) (j - k) (j^2 - 3 k j + k^2) j$$

$$R_3 := (j - 2 k) (j^2 + k j - k^2) k j$$

$$R_4 := - (j - k) (j^2 - k j - k^2) (-k + 2 j) k$$

$$R_5 := - (j - k) (-2 k j^3 + j^4 - j^2 k^2 + k^4)$$

$$R_6 := (j^4 - 4 k j^3 + j^2 k^2 + 2 k^3 j - k^4) k$$

$$R_7 := (j^4 - 4 k j^3 + 5 j^2 k^2 - k^4) j$$

On expansion

$$\begin{aligned}
 F_7 = & 2 j^3 k^3 (-k + 2j) (j - 2k) (j + k) \\
 & (j^2 + kj - k^2) (j^2 - kj - k^2) (j^2 - 3kj + k^2) \\
 & (-3j^2k + k^3 + j^3) (j^4 - 4kj^3 + 5j^2k^2 - k^4) \\
 & (-2kj^3 + j^4 - j^2k^2 + k^4) (j - k)^3 \\
 & (j^4 - 4kj^3 + j^2k^2 + 2k^3j - k^4) (j^2 - kj + k^2)
 \end{aligned}$$

For example with  $j := 2$  and  $k := 3$

$$\begin{aligned}
 & (t - 7) (t - 50) (t + 24) (t + 33) \\
 & (t - 13) (t + 51) (t - 38) \\
 & - (t + 7) (t + 50) (t - 24) (t - 33) \\
 & (t + 13) (t - 51) (t + 38) \\
 & = 13967553600
 \end{aligned}$$

**Size 8.** A (homogenous) size 8 solution due to Chernick

$$F_8 := (t^2 - R_1^2) (t^2 - R_2^2) (t^2 - R_3^2) (t^2 - R_4^2) \\ - (t^2 - R_5^2) (t^2 - R_6^2) (t^2 - R_7^2) (t^2 - R_8^2)$$

where

$$R_1 := 5m^2 + 9mn + 10n^2$$

$$R_2 := m^2 - 13mn - 6n^2$$

$$R_3 := 7m^2 - 5mn - 8n^2$$

$$R_4 := 9m^2 + 7mn - 4n^2$$

$$R_5 := 9m^2 + 5mn + 4n^2$$

$$R_6 := m^2 + 15mn + 8n^2$$

$$R_7 := 5m^2 - 7mn - 10n^2$$

$$R_8 := 7m^2 + 5mn - 6n^2$$

**Size 9.** We know no parametric solution of size 9. Indeed only two solutions are known. Both are symmetric and they are the following

$$[-98, -82, -58, -34, 13, 16, 69, 75, 99]$$

and

$$[174, 148, 132, 50, 8, -63, -119, -161, -169]$$

**Size 10.** The following size 10 example is due to Letac (and Smyth )

$$F_{10} :=$$

$$\begin{aligned} & \left( t^2 - R_1^2 \right) \left( t^2 - R_2^2 \right) \left( t^2 - R_3^2 \right) \\ & \left( t^2 - R_4^2 \right) \left( t^2 - R_5^2 \right) \\ & - \left( t^2 - R_6^2 \right) \left( t^2 - R_7^2 \right) \left( t^2 - R_8^2 \right) \\ & \left( t^2 - R_9^2 \right) \left( t^2 - R_{10}^2 \right) \end{aligned}$$

where

$$R_1 := (4n + 4m)$$

$$R_2 := (mn + n + m - 11)$$

$$R_3 := (mn - n - m - 11)$$

$$R_4 := (mn + 3n - 3m + 11)$$

$$R_5 := (mn - 3n + 3m + 11)$$

$$R_6 := (4n - 4m)$$

$$R_7 := (-mn + n - m - 11)$$

$$R_8 := (-mn - n + m - 11)$$

$$R_9 := (-mn + 3n + 3m + 11)$$

$$R_{10} := (-mn - 3n - 3m + 11)$$

On expansion

$$F_{10} := c_0 + c_2 t^2 + c_4 t^4 + c_6 t^6$$

And each coefficient except  $c_0$  has a factor

$$m^2 n^2 - 13n^2 + 121 - 13m^2$$

So any solution of the above biquadratic gives a size 10 solution. For example:

$$n := 153/61 \text{ and } m = 191/79$$

$$n := -296313/249661 \text{ and } m = -1264969/424999$$