

MATHEMATICS 152 98-2 Solutions for Assignment 5

$$2. \int x \cos x \, dx = x \sin x - \int \sin x \, dx = x \sin x + \cos x + C.$$

$$4. \int x \ln x \, dx = \frac{1}{2} x^2 \ln x - \int \frac{1}{2} x \, dx = \frac{1}{2} x^2 \ln x - \frac{1}{4} x^2 + C.$$

$$10. \int \theta \sec^2 \theta \, d\theta = \theta \tan \theta - \int \tan \theta \, d\theta = \theta \tan \theta - \int \frac{\sin \theta}{\cos \theta} \, d\theta = \theta \tan \theta + \ln |\cos \theta| + C.$$

$$12. \int t^3 e^t \, dt = t^3 e^t - \int 3t^2 e^t \, dt = t^3 e^t - 3t^2 e^t + \int 6t e^t \, dt = t^3 e^t - 3t^2 e^t + 6t e^t - \int 6 e^t \, dt = t^3 e^t - 3t^2 e^t + 6t e^t - 6e^t + C = (t^3 - 3t^2 + 6t - 6)e^t + C.$$

(You could do something similar for $\int P(t)e^t \, dt$, where $P(t)$ is any polynomial; eventually you would differentiate the polynomial to death.)

$$18. \int_1^4 \sqrt{t} \ln t \, dt = \frac{2}{3} t^{3/2} \ln t \Big|_1^4 - \int_1^4 \frac{2}{3} t^{1/2} \, dt = \frac{16}{3} \ln 4 - \frac{4}{9} t^{3/2} \Big|_1^4 = \frac{16}{3} \ln 4 - \frac{28}{9}.$$

$$24. \int x^3 e^{x^2} \, dx = \int \left(\frac{1}{2} x^2 \right) \cdot (2x e^{x^2}) \, dx = \frac{1}{2} x^2 e^{x^2} - \int x e^{x^2} \, dx = \frac{1}{2} x^2 e^{x^2} - \frac{1}{2} e^{x^2} + C = \frac{1}{2} (x^2 - 1) e^{x^2} + C.$$

$$30. \int \sin(\ln x) \, dx = \int x \sin(\ln x) \cdot \frac{1}{x} \, dx = -x \cos(\ln x) + \int \cos(\ln x) \, dx = -x \cos(\ln x) + \int x \cos(\ln x) \cdot \frac{1}{x} \, dx = -x \cos(\ln x) + x \sin(\ln x) - \int \sin(\ln x) \, dx.$$

So $2 \int \sin(\ln x) \, dx = -x \cos(\ln x) + x \sin(\ln x) + K$, and

$$\int \sin(\ln x) \, dx = -\frac{1}{2} x \cos(\ln x) + \frac{1}{2} x \sin(\ln x) + C = \frac{x}{2} [\sin(\ln x) - \cos(\ln x)] + C.$$

$$38. (a) \int \cos^n x \, dx = \int \cos^{n-1} x \cos x \, dx = \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x \sin^2 x \, dx = \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x (1 - \cos^2 x) \, dx = \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x \, dx - (n-1) \int \cos^n x \, dx.$$

So $n \int \cos^n x \, dx = \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x \, dx$, and

$$\int \cos^n x \, dx = \frac{1}{n} \cos^{n-1} x \sin x + \frac{n-1}{n} \int \cos^{n-2} x \, dx.$$

$$\begin{aligned} \text{(b)} \quad \int \cos^2 x \, dx &= \frac{1}{2} \cos x \sin x + \frac{1}{2} \int 1 \, dx = \frac{1}{2} \cos x \sin x + \frac{1}{2} x + C = \\ &= \frac{1}{4} \sin(2x) + \frac{1}{2} x + C. \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \int \cos^4 x \, dx &= \frac{1}{4} \cos^3 x \sin x + \frac{3}{4} \int \cos^2 x \, dx = \\ &= \frac{1}{4} \cos^3 x \sin x + \frac{3}{4} \left(\frac{1}{2} \cos x \sin x + \frac{1}{2} x \right) + C = \\ &= \frac{1}{4} \cos^3 x \sin x + \frac{3}{8} \cos x \sin x + \frac{3}{8} x + C = \frac{1}{4} \sin(2x) + \frac{1}{32} \sin(4x) + \frac{3}{8} x + C. \end{aligned}$$

$$42. \quad \int x^n e^x \, dx = x^n e^x - \int n x^{n-1} e^x \, dx = x^n e^x - n \int x^{n-1} e^x \, dx.$$


$$\begin{aligned} 46. \quad \int x^4 e^x \, dx &= x^4 e^x - 4 \int x^3 e^x \, dx = x^4 e^x - 4x^3 e^x + 12 \int x^2 e^x \, dx = \\ &= x^4 e^x - 4x^3 e^x + 12x^2 e^x - 24 \int x e^x \, dx = x^4 e^x - 4x^3 e^x + 12x^2 e^x - 24x e^x + 24 \int e^x \, dx = \\ &= x^4 e^x - 4x^3 e^x + 12x^2 e^x - 24x e^x + 24e^x + C = (x^4 - 4x^3 + 12x^2 - 24x + 24) e^x + C. \end{aligned}$$

48. $y = 5 \ln x$ meets $y = x \ln x$ at the point $(1, 0)$, where $\ln x = 0$, and also at the point $(5, 5 \ln 5)$, where $x = 5$. Between these points $\ln x > 0$ and $x < 5$, so $y = x \ln x$ is the lower curve and $y = 5 \ln x$ is the upper curve.

The enclosed area is given by

$$\begin{aligned} \int_1^5 (5 \ln x - x \ln x) \, dx &= \int_1^5 (5 - x) \ln x \, dx = \\ &= -\frac{1}{2} (5 - x)^2 \ln x \Big|_1^5 + \frac{1}{2} \int_1^5 \frac{(5 - x)^2}{x} \, dx = \\ &= 0 + \frac{1}{2} \int_1^5 \left(\frac{25}{x} - 10 + x \right) \, dx = \\ &= \frac{1}{2} \left(25 \ln x - 10x + \frac{1}{2} x^2 \right) \Big|_1^5 = \\ &= \frac{1}{2} (25 \ln 5 - 28) = 12.5 \ln 5 - 14. \end{aligned}$$

See graph above and to the right.



For Exercise 48

$$2. \int_0^{\pi/2} \cos^2 x \, dx = \left. \frac{1}{2} \cos x \sin x \right|_0^{\pi/2} + \int_0^{\pi/2} \frac{1}{2} \, dx = 0 + \left. \frac{1}{2} x \right|_0^{\pi/2} = \frac{\pi}{4}.$$

$$\text{Alternatively } \int_0^{\pi/2} \cos^2 x \, dx = \int_0^{\pi/2} \frac{1 + \cos(2x)}{2} \, dx = \left. \left(\frac{1}{2} x + \frac{1}{4} \sin(2x) \right) \right|_0^{\pi/2} = \frac{\pi}{4}.$$

Alternatively (and the laziest way of all) if $u = \frac{\pi}{2} - x$ so that $du = -dx$, then

$$\int_0^{\pi/2} \cos^2 x \, dx = -\int_{\pi/2}^0 \sin^2 u \, du = \int_0^{\pi/2} \sin^2 u \, du.$$

The name of the variable of integration is irrelevant, so $\int_0^{\pi/2} \sin^2 u \, du = \int_0^{\pi/2} \sin^2 x \, dx$,

$$\text{and thus } \int_0^{\pi/2} \cos^2 x \, dx = \int_0^{\pi/2} \sin^2 u \, du = \int_0^{\pi/2} \sin^2 x \, dx = \int_0^{\pi/2} [1 - \cos^2 x] \, dx.$$

$$\text{Consequently } 2 \int_0^{\pi/2} \cos^2 x \, dx = \int_0^{\pi/2} 1 \, dx = x \Big|_0^{\pi/2} = \frac{\pi}{2}, \text{ and } \int_0^{\pi/2} \cos^2 x \, dx = \frac{\pi}{4}.$$

$$6. \int \sin^4 x \cos^3 x \, dx = \int \sin^4 x (1 - \sin^2 x) \cos x \, dx = \int (\sin^4 x - \sin^6 x) \cos x \, dx = \\ = \frac{1}{5} \sin^5 x - \frac{1}{7} \sin^7 x + C.$$

$$10. \int \sin \left(x + \frac{\pi}{6} \right) \cos x \, dx = \int \frac{1}{2} \left[\sin \frac{\pi}{6} + \sin \left(2x + \frac{\pi}{6} \right) \right] dx = \\ = \left(\frac{1}{2} \sin \frac{\pi}{6} \right) \cdot x - \frac{1}{4} \cos \left(2x + \frac{\pi}{6} \right) + C = \frac{1}{4} x - \frac{1}{4} \cos \left(2x + \frac{\pi}{6} \right) + C.$$

$$\text{Alternatively } \int \sin \left(x + \frac{\pi}{6} \right) \cos x \, dx = \int \left[\sin x \cos \frac{\pi}{6} + \cos x \sin \frac{\pi}{6} \right] \cos x \, dx = \\ = \int \left(\frac{\sqrt{3}}{2} \sin x \cos x + \frac{1}{2} \cos^2 x \right) dx = \int \left(\frac{\sqrt{3}}{4} \sin(2x) + \frac{1 + \cos(2x)}{4} \right) dx = \\ = -\frac{\sqrt{3}}{8} \cos(2x) + \frac{1}{4} x + \frac{1}{8} \sin(2x) + C. \text{ These answers really are the same.}$$

$$14. \int x \sin^3(x^2) \, dx = \int x \sin^2(x^2) \sin(x^2) \, dx = \int x [1 - \cos^2(x^2)] \sin(x^2) \, dx = \\ = \int [(\sin(x^2) \cdot x - \cos^2(x^2) \sin(x^2) \cdot x)] \, dx = -\frac{1}{2} \cos(x^2) + \frac{1}{6} \cos^3(x^2) + C.$$

$$26. \int \tan^3 x \sec^3 x \, dx = \int \tan^2 x \sec^2 x (\sec x \tan x) \, dx = \\ = \int [\sec^2 x - 1] \sec^2 x (\sec x \tan x) \, dx = \int [\sec^4 x - \sec^2 x] (\sec x \tan x) \, dx = \\ = \frac{1}{5} \sec^5 x - \frac{1}{3} \sec^3 x + C.$$

$$36. \int \cot^3 x \csc^4 x \, dx = \int \cot^3 x [\cot^2 x + 1] \csc^2 x \, dx = \int [\cot^5 x + \cot^3 x] \csc^2 x \, dx = \\ = -\frac{1}{6} \cot^6 x - \frac{1}{4} \cot^4 x + C.$$

$$42. \int \sin(3x) \sin(6x) \, dx = \int \sin(3x) [2 \sin(3x) \cos(3x)] \, dx = \int 2 \sin^2(3x) \cos(3x) \, dx = \\ = \frac{2}{9} \sin^3(3x) + C.$$

$$\text{Alternatively } \int \sin(3x) \sin(6x) \, dx = \int \frac{1}{2} [\cos(3x) - \cos(9x)] \, dx = \\ = \frac{1}{6} \sin(3x) - \frac{1}{18} \sin(9x) + C. \text{ These answers really are the same.}$$

$$48. (a) \int \sin x \cos x \, dx = \int (-\cos x) \left(\frac{d}{dx} \cos x \right) \, dx = -\frac{1}{2} \cos^2 x + C_1.$$

$$(b) \int \sin x \cos x \, dx = \int \sin x \left(\frac{d}{dx} \sin x \right) \, dx = \frac{1}{2} \sin^2 x + C_2.$$

$$(c) \int \sin x \cos x \, dx = \int \frac{1}{2} \sin(2x) \, dx = -\frac{1}{4} \cos(2x) + C_3.$$

$$(d) \int \sin x \cos x \, dx = -\cos x \cdot \cos x - \int \cos x \cdot \sin x \, dx, \text{ hence}$$

$$2 \int \sin x \cos x \, dx = -\cos^2 x + K_1 \text{ and } \int \sin x \cos x \, dx = -\frac{1}{2} \cos^2 x + C_1.$$

$$\text{Alternatively } \int \sin x \cos x \, dx = \sin x \cdot \sin x - \int \cos x \cdot \sin x \, dx, \text{ hence}$$

$$2 \int \sin x \cos x \, dx = \sin^2 x + K_2 \text{ and } \int \sin x \cos x \, dx = \frac{1}{2} \sin^2 x + C_2.$$

$$\text{Here } C_2 = C_1 - \frac{1}{2} \text{ and } C_3 = C_1 - \frac{1}{4}.$$


50. In the region $0 \leq x \leq \frac{\pi}{2}$, the curves $y = \sin x$ and $y = 2 \sin^2 x$ meet at $(0, 0)$ and at $(\pi/6, 1/2)$, since $\sin 0 = 0$ and $2 \sin(\pi/6) = 1$.

For $0 < x < \frac{\pi}{6}$, $y = 2 \sin^2 x$ is the lower curve; for $\frac{\pi}{6} < x \leq \frac{\pi}{2}$, $y = \sin x$ is the lower curve. Thus the height of the region is $\sin x - 2 \sin^2 x$ for $0 \leq x \leq \frac{\pi}{6}$.

The height is $2 \sin^2 x - \sin x$ for $\frac{\pi}{6} \leq x \leq \frac{\pi}{2}$.

$$\text{The area is } \int_0^{\pi/6} [\sin x - 2 \sin^2 x] \, dx + \int_{\pi/6}^{\pi/2} [2 \sin^2 x - \sin x] \, dx = \\ = \int_0^{\pi/6} [\sin x + \cos(2x) - 1] \, dx + \int_{\pi/6}^{\pi/2} [-\sin x - \cos(2x) + 1] \, dx = \\ = \left(-\cos x + \frac{1}{2} \sin(2x) - x \right) \Big|_0^{\pi/6} + \left(\cos x - \frac{1}{2} \sin(2x) + x \right) \Big|_{\pi/6}^{\pi/2} = 1 - \frac{\sqrt{3}}{2} + \frac{\pi}{6}.$$


See graph above and to the right.



56. Using the method of washers, the volume is

$$\begin{aligned} V &= \pi \int_0^{\pi/2} [(1 - 0)^2 - (1 - \cos x)^2] dx = \\ &= \pi \int_0^{\pi/2} [2 \cos x - \cos^2 x] dx = \\ &= \pi \int_0^{\pi/2} \left(2 \cos x - \frac{1 + \cos(2x)}{2} \right) dx = \\ &= \pi \left(2 \sin x - \frac{1}{2} x - \frac{1}{4} \sin(2x) \right) \Big|_0^{\pi/2} = 2\pi - \frac{\pi^2}{4}. \end{aligned}$$

The method of cylindrical shells can also be used, but it involves integrating $(1 - y) \cos^{-1} y$. That integration (by parts) is somewhat tricky.



For Exercise 56