

Rational Approximation with Varying Weights I

P. Borwein*, E. A. Rakhmanov[†] and E. B. Saff[‡]

Abstract

We investigate two problems concerning uniform approximation by weighted rationals $\{w^n r_n\}_{n=1}^\infty$, where $r_n = p_n/q_n$ is a rational function of order n . Namely, for $w(x) := e^x$ we prove that uniform convergence to 1 of $w^n r_n$ is not possible on any interval $[0, a]$ with $a > 2\pi$. For $w(x) := x^\theta$, $\theta > 1$, we show that uniform convergence to 1 of $w^n r_n$ is not possible on any interval $[b, 1]$ with $b < \tan^4(\pi(\theta - 1)/4\theta)$. (The latter result can be interpreted as a rational analogue of results concerning “incomplete polynomials”.) More generally, for $\alpha \geq 0$, $\beta \geq 0$, $\alpha + \beta > 0$, we investigate for $w(x) = e^x$ and $w(x) = x^\theta$, the possibility of approximation by $\{w^n p_n/q_n\}_{n=1}^\infty$, where $\deg p_n \leq \alpha n$, $\deg q_n \leq \beta n$. The analysis utilizes potential theoretic

*Research supported, in part, by NSERC of Canada.

[†]The research of this author was conducted while visiting the University of South Florida, Tampa.

[‡]The research of this author was supported, in part, by the U.S. National Science Foundation under grant DMS 920-3659.

methods. These are essentially sharp results though this will not be established in this paper.

1 Introduction and Main Results

1.1 . For a positive, continuous function $f(x)$ on $\mathbf{R}_+ = [0, +\infty)$ we define

$$\delta_n(f; R) := \inf_{r \in \mathcal{R}_n} \left\| \frac{r(x)}{f(x)} - 1 \right\|_{[0, R]}, \quad (1.1)$$

where \mathcal{R}_n is the set of all real rational functions of order $\leq n$ and $\| \cdot \|_{[a, b]}$ denotes the sup norm over the interval $[a, b]$. That is, we consider the best relative rational approximations to f on $[0, R]$. It is clear that $\delta_n(f; R) \rightarrow 0$ as $n \rightarrow \infty$ for any fixed $R \in (0, +\infty)$ and, moreover, it is always possible to find an increasing sequence $R_n \rightarrow \infty$ satisfying the condition

$$\delta_n(f, R_n) \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (1.2)$$

On the other hand if $f(x)$, say, decreases as $x \rightarrow \infty$, then R_n satisfying (??) cannot increase arbitrarily fast, which raises the question about the maximum possible rate of increase of R_n .

Here we consider this question for the ‘‘model’’ function $f(x) = e^{-x}$.

Theorem 1.1 *If (??) is true for $f(x) = e^{-x}$, then*

$$R_n \leq (1 + \epsilon)2\pi n$$

for any $\epsilon > 0$ and $n \geq n(\epsilon)$.

Remark A. We do not know whether the inequality $R_n \leq 2\pi n$ is true for large enough n . However, Theorem ?? is sharp in the sense that the constant 2π cannot be replaced by any smaller constant. The proof of the last assertion, which is substantial, will appear in a future paper (it is based on constructing rational approximations following the method developed in [GR]).

Remark B. A weaker version of Theorem 1.1 with a constant of 8 replacing 2π follows easily from a result in [B] which says that, for a non-zero rational function of order $\leq n$,

$$m \left\{ x \in \mathbf{R} : \frac{r'_n(x)}{r_n(x)} \geq n \right\} \leq 8.$$

Here m denotes Lebesgue measure. It is reasonable to hypothesize from the results of this paper and its sequel that the constant in the above inequality should be 2π .

In Sec. 1.3 below we present a generalization of Theorem ?? concerning approximation by ray sequences of rational functions.

We note that the corresponding question about relative *polynomial* approximation is important for the investigation of strong asymptotics for orthogonal polynomials on \mathbf{R} and \mathbf{R}_+ . Such results dealing with relative polynomial approximation were obtained in [LS], [LR], [ST], [To].

1.2 . Another problem considered in this paper is the approximation of the sequence $x^{n\theta}$ on subintervals of $[0, 1]$. For $\theta > 0$, we set

$$\Delta_n(\theta, b) := \inf_{r \in \mathcal{R}_n} \|x^{n\theta} r(x) - 1\|_{[b, 1]}, \quad b \in (0, 1). \quad (1.3)$$

Theorem 1.2 *If $\Delta_n(\theta, b) \rightarrow 0$ as $n \rightarrow \infty$ and $\theta > 1$, then*

$$b \geq \tan^4 \left(\frac{\pi \theta - 1}{4 \theta} \right). \quad (1.4)$$

Remark C. It is clear that for $\theta \leq 1$ approximation is possible over $[0, 1]$. Furthermore, if $\theta > 1$, the right-hand side of (??) cannot be replaced by any larger constant (again this fact will appear in a future paper).

In Sec. 1.4 we present a more general result dealing with approximation by ray sequences of rationals.

We note that Theorem ?? is closely related to the completeness of the system of “incomplete rational functions”

$$\left\{ x^{n\theta} \frac{p_n(x)}{q_n(x)} : \deg p_n, \deg q_n \leq n \right\}$$

in $C[b, 1]$. If $n\theta$ is an integer, then $x^{n\theta} p_n(x)/q_n(x)$ may be interpreted as a rational function of order $n(1+\theta)$ with $n\theta$ poles fixed at ∞ and $n\theta$ zeros fixed at 0. Corresponding questions for incomplete polynomials are considered in [Lo], [SV]. Related questions for incomplete rationals in the complex plane are treated in [BC].

1.3 . For fixed $\alpha, \beta \geq 0$, $\alpha + \beta > 0$, set

$$\delta_n(a; \alpha, \beta) := \inf_{p, q} \left\| e^{nx} \frac{p(x)}{q(x)} - 1 \right\|_{[0, a]}, \quad (1.5)$$

where the infimum is taken over all polynomials p, q with $\deg p \leq \alpha n$, $\deg q \leq \beta n$. We also define

$$a^* := a^*(\alpha, \beta) := \sup \{ a : \delta_n(a; \alpha, \beta) \rightarrow 0 \text{ as } n \rightarrow \infty \}. \quad (1.6)$$

Theorem 1.3 *We have the inequality $a^* \leq \hat{a}$, where $\hat{a} := 2\pi\alpha$ for $\alpha = \beta$ and*

$$\hat{a} := \frac{2(\alpha - \beta)}{1 - 2\hat{y}} \quad (1.7)$$

where $\hat{y} = \hat{y}(\alpha, \beta)$ is the root of the equation

$$g(y) := \frac{\sqrt{y(1-y)}}{1-2y} - \sin^{-1} \sqrt{y} = \frac{\pi}{2} \frac{\beta}{\alpha - \beta} \quad (1.8)$$

for $\alpha \neq \beta$.

Remark D. It can be shown that $\hat{a}(\alpha, \beta) \rightarrow 2\pi\alpha$ as $\beta \rightarrow \alpha$ (for fixed α), so the function $\hat{a}(\alpha, \beta)$ is continuous.

Remark E. Both functions $a^*(\alpha, \beta)$ and $\hat{a}(\alpha, \beta)$ are symmetric and therefore we need only consider the case $\alpha \geq \beta$ (the symmetry of a^* can be seen on making the change of variables $x \rightarrow a - x$; the symmetry of \hat{a} follows from the identity $g(1-y) + g(y) = -\pi/2$). The case $\alpha \geq \beta$ is equivalent to $\hat{y} \in [0, 1/2]$. In this interval we have

$$g(y) = 2 \int_0^y \frac{\sqrt{t(1-t)}}{(1-2t)^2} dt, \quad (1.9)$$

and therefore $g(y)$ is increasing from 0 to ∞ on $[0, 1/2]$. Hence the equation $g(y) = (\pi/2) \beta / (\alpha - \beta)$ has a unique root for any $\alpha \geq \beta \geq 0$ ($\alpha + \beta > 0$).

Remark F. For $\alpha + \beta = 1$ (i.e., for a fixed number of free parameters in p_n/q_n) the function $\hat{a}(\alpha, \beta) = \hat{a}(1 - \beta, \beta)$ takes its maximum value over $\beta \in [0, 1]$ at $\beta = 1/2$. This means that diagonal approximations ($\alpha = \beta$)

are the most effective among all ray sequences with the same number of free parameters.

1.4 . For fixed $\theta > 0$, $\alpha, \beta \geq 0$, $\alpha + \beta > 0$, we define

$$\Delta_n(b, \theta; \alpha, \beta) := \inf_{p, q} \left\| x^{n\theta} \frac{p(x)}{q(x)} - 1 \right\|_{[b, 1]}, \quad (1.10)$$

where the infimum is taken over all polynomials p, q satisfying $\deg p \leq \alpha n$, $\deg q \leq \beta n$, and we set

$$b^* = b^*(\theta; \alpha, \beta) := \inf \{b : \Delta_n(b, \theta; \alpha, \beta) \rightarrow 0 \text{ as } n \rightarrow \infty\}. \quad (1.11)$$

Theorem 1.4 *We have the estimate $b^* \geq \hat{b}$, where $\hat{b} = \hat{b}(\theta; \alpha, \beta)$ is the unique root of the equation*

$$f(b) := \frac{1}{\pi} \int_0^b \frac{\sqrt{(\xi - \sqrt{t})(1 - \xi\sqrt{t})}}{t^{3/4}(1-t)} dt = 1 - \frac{\beta}{\theta}; \quad \xi := 1 + \frac{\alpha}{\theta} - \frac{\beta}{\theta}, \quad (1.12)$$

when $\beta/\theta \leq 1$ and $\hat{b} := 0$ when $\beta/\theta > 1$.

We note that for $\beta/\theta > 1$ the fact that $b^* = 0$ is easily seen.

We shall also obtain the following representation for f :

$$f(b) = 1 - \frac{2}{\pi} \sin^{-1} \sqrt{\frac{1 - \xi\sqrt{b}}{1 - b}} + \xi \frac{2}{\pi} \sin^{-1} \sqrt{\frac{\sqrt{b} (1 - \xi\sqrt{b})}{\xi (1 - b)}}, \quad (1.13)$$

when $\sqrt{b} \leq \xi \leq 1/\sqrt{b}$. (We shall see that \hat{b} satisfies these inequalities.) The two important particular cases $\alpha = 0$ and $\beta = 0$ have already been considered in [SV], [G], [BS] and the following results obtained:

If $\beta = 0$, then $b^* = (1 + \alpha/\theta)^{-2}$.

If $\alpha = 0$, then $b^* = (1 - \beta/\theta)^2$ for $\beta/\theta < 1$ and $b^* = 0$ if $\beta/\theta \geq 1$.

Note that the corresponding lower estimates are included in Theorem ??.

2 Proofs of Theorems 1.1 and 1.3

We denote by $V(x, \mu)$ the logarithmic potential for the measure $d\mu$:

$$V(x, \mu) := \int \log \frac{1}{|x - t|} d\mu(t).$$

We fix $a > 0$ and define the two distributions:

$$\sigma_1(t) := \frac{1}{\pi} \sqrt{\frac{a-t}{t}}, \quad t \in [0, a], \quad (2.1)$$

$$\sigma_0(t) := \frac{1}{\pi} \frac{1}{\sqrt{t(a-t)}}, \quad t \in (0, a). \quad (2.1a)$$

The following properties of the corresponding logarithmic potentials are easily verified (see Appendix):

$$V(x, \sigma_1 dt) = -x + \text{const}, \quad x \in [0, a], \quad (2.2)$$

$$V(x, \sigma_0 dt) = \log(4/a), \quad x \in [0, a]. \quad (2.3)$$

For $x \in \mathbf{R}$ we define the function

$$\sigma(t, x) := \sigma_1(t) - x\sigma_0(t), \quad 0 \leq t \leq a. \quad (2.4)$$

For each fixed x , let $\sigma(t, x) = \sigma^+(t, x) - \sigma^-(t, x)$ be the Jordan decomposition of the measure $\sigma(t, x)dt$ in $[0, a]$ and set

$$p(x) := p(x, a) := \int \sigma^+(t, x)dt \quad (2.5)$$

$$n(x) := n(x, a) := \int \sigma^-(t, x)dt. \quad (2.6)$$

Lemma 2.1 *With the notation of Theorem ?? we have the following implication: if $a = a(\alpha, \beta) < a^* = a^*(\alpha, \beta)$, then there exists an $x \in \mathbf{R}$ such that*

$$p(x, a) \leq \beta \quad \text{and} \quad n(x, a) \leq \alpha.$$

Proof. The condition $a < a^*$ means that (α, β) -approximation to e^{nx} on $[0, a]$ is possible. In other words, there exist two sequences of polynomials

$$p_n \in \mathcal{P}_{[\alpha n]}, \quad q_n \in \mathcal{P}_{[\beta n]}, \quad (2.7)$$

with

$$\delta_n := \left\| e^{nx} \frac{p_n(x)}{q_n(x)} - 1 \right\|_{[0, a]} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (2.8)$$

Set

$$\chi_{n,\alpha} := \frac{1}{n} \sum_{p_n(z)=0} \delta(z), \quad \chi_{n,\beta} := \frac{1}{n} \sum_{q_n(z)=0} \delta(z),$$

where $\delta(z)$ denotes the unit point measure at z , and define $\mu_{n,\alpha}$ and $\mu_{n,\beta}$ to be the balayage (cf. [La]) of $\chi_{n,\alpha}$ and $\chi_{n,\beta}$ into $[0, a]$, respectively. Then we have (cf. [La])

$$\frac{1}{n} \log \frac{1}{|p_n(x)|} = V(x, \mu_{n,\alpha}) + \omega_{n,\alpha}, \quad x \in [0, a], \quad (2.9)$$

$$\frac{1}{n} \log \frac{1}{|q_n(x)|} = V(x, \mu_{n,\beta}) + \omega_{n,\beta}, \quad x \in [0, a], \quad (2.10)$$

where $\omega_{n,\alpha}$ and $\omega_{n,\beta}$ are constants depending on n . It follows from these two representations and (2.2) that

$$\frac{1}{n} \log \left| e^{nx} \frac{p_n(x)}{q_n(x)} \right| = V(x, \mu_{n,\beta} - \mu_{n,\alpha} - \sigma_1 dt) + \omega_n, \quad x \in [0, a], \quad (2.11)$$

where ω_n is a constant.

From (??), we deduce that

$$\left| \frac{1}{n} \log \left(e^{nx} \frac{p_n(x)}{q_n(x)} \right) \right| \leq \frac{1}{n} |\log(1 - \delta_n)| \rightarrow 0$$

uniformly on $[0, a]$ as $n \rightarrow \infty$. Therefore, with

$$\mu_n := \mu_{n,\beta} - \mu_{n,\alpha} - \sigma_1 dt$$

we have, uniformly on $[0, a]$,

$$V(x, \mu_n) + \omega_n \rightarrow 0. \quad (2.12)$$

Furthermore, we see from (??) that

$$\begin{aligned} \|\mu_{n,\alpha}\| &= \|\chi_{n,\alpha}\| = \frac{1}{n} \deg p_n \leq \alpha, \\ \|\mu_{n,\beta}\| &= \|\chi_{n,\beta}\| = \frac{1}{n} \deg q_n \leq \beta. \end{aligned}$$

Hence, we can find a subsequence $\Lambda \subset \mathbf{N}$ and positive measures μ_α, μ_β such that as $n \rightarrow \infty$, $n \in \Lambda$,

$$\mu_{n,\alpha} \xrightarrow{*} \mu_\alpha, \quad \|\mu_\alpha\| \leq \alpha; \quad \mu_{n,\beta} \xrightarrow{*} \mu_\beta, \quad \|\mu_\beta\| \leq \beta; \quad (2.13)$$

where $\xrightarrow{*}$ denotes weak-star convergence.

It follows from (??) that $\mu_n \xrightarrow{*} \mu := \mu_\beta - \mu_\alpha - \sigma_1 dt$ and therefore as $n \rightarrow \infty$, $n \in \Lambda$,

$$\mu_n(\mathbf{C}) \rightarrow \mu(\mathbf{C}), \quad V(z, \mu_n) \rightarrow V(z, \mu), \quad z \in \mathbf{C} \setminus [0, a]. \quad (2.14)$$

Furthermore, on integrating (??) with respect to the (unit) equilibrium measure $\sigma_0(x)dx$ and utilizing (??) we obtain

$$\begin{aligned} \int_0^a V(x, \mu_n) \sigma_0(x) dx + \omega_n &= \int_0^a V(t, \sigma_0 dx) d\mu_n(t) + \omega_n \\ &= \mu_n(\mathbf{C}) \log(4/a) + \omega_n \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$, $n \in \Lambda$, and so from (??) we have

$$\lim_{\substack{n \rightarrow \infty \\ n \in \Lambda}} \omega_n = -\mu(\mathbf{C}) \log(4/a). \quad (2.15)$$

Next observe that for n sufficiently large, $p_n(x)$ and $q_n(x)$ do not vanish on $[0, a]$; hence from (??) and (??) it follows that $V(x, \mu_n)$ is finite and continuous on $\text{supp}(\mu_n)$ and therefore $V(z, \mu_n)$ is continuous on \mathbf{C} . Consequently,

$$h_n(z) := V(z, \mu_n) - \mu_n(\mathbf{C})V(z, \sigma_0 dt)$$

is continuous on $\overline{\mathbf{C}} = \mathbf{C} \cup \{\infty\}$ and harmonic in $\mathbf{C} \setminus [0, a]$. Thus, by (??), (??), (??) and the maximum principle, we have

$$\lim_{\substack{n \rightarrow \infty \\ n \in \Lambda}} h_n(z) = 0, \quad z \in \overline{\mathbf{C}}.$$

On the other hand, (??) yields

$$\lim_{\substack{n \rightarrow \infty \\ n \in \Lambda}} h_n(z) = V(z, \mu) - \mu(\mathbf{C})V(z, \sigma_0 dt), \quad z \in \mathbf{C} \setminus [0, a],$$

and so

$$V(z, \mu) = V(z, \lambda \sigma_0 dt), \quad z \in \mathbf{C} \setminus [0, a],$$

where $\lambda := \mu(\mathbf{C})$. Since the potential of the signed measure $\mu - \lambda \sigma_0 dt$ vanishes outside a set of 2-dimensional Lebesgue measure zero, we have $\mu = \lambda \sigma_0 dt$

and we obtain

$$\mu_\beta - \mu_\alpha = \sigma_1 dt - \lambda \sigma_0 dt = \sigma(t, \lambda) dt$$

(see (??)). Due to the minimizing property of the Jordan decomposition we then have

$$\alpha \geq \|\mu_\alpha\| \geq \|\sigma^-(t, \lambda)dt\| = n(\lambda, a),$$

$$\beta \geq \|\mu_\beta\| \geq \|\sigma^+(t, \lambda)dt\| = p(\lambda, a),$$

which completes the proof. ■

Lemma 2.2 *The following properties of the functions $p(x) := p(x, a)$ and $n(x) := n(x, a)$ (cf. (??) and (??)) hold for any fixed $a > 0$:*

$$(i) \quad p(x) - n(x) = \frac{a}{2} - x, \quad x \in \mathbf{R};$$

$$(ii) \quad p(x) = \frac{a}{2} - x \text{ and } n(x) = 0 \text{ for } x \leq 0,$$

$$p(x) = 0 \text{ and } n(x) = x - \frac{a}{2} \text{ for } x \geq a;$$

$$(iii) \quad p(x) = \frac{a}{\pi} \left\{ \left(1 - 2\frac{x}{a}\right) \sin^{-1} \sqrt{1 - \frac{x}{a}} + \sqrt{\frac{x}{a} \left(1 - \frac{x}{a}\right)} \right\}, \quad x \in [0, a];$$

$$(iv) \quad p'(x) = -\frac{2}{\pi} \sin^{-1} \sqrt{1 - \frac{x}{a}}, \quad x \in [0, a],$$

$$n'(x) = \frac{2}{\pi} \cos^{-1} \sqrt{1 - \frac{x}{a}}, \quad x \in [0, a].$$

Proof. We have from definitions (??), (??) and (??)–(??)

$$p(x) - n(x) = \int_0^a \sigma(t, x)dt = \frac{1}{\pi} \int_0^a \sqrt{\frac{a-t}{t}} dt - x \frac{1}{\pi} \int_0^a \frac{dt}{\sqrt{t(a-t)}} = \frac{a}{2} - x.$$

Furthermore, we have

$$\sigma(t, x) = \frac{1}{\pi} \frac{a - t - x}{\sqrt{t(a - t)}} \geq 0 \text{ for } t \in [0, a], x \leq 0,$$

and therefore $n(x) = 0$ for $x \leq 0$. Likewise $\sigma(t, x) \leq 0$ for $t \in [0, a]$ and $x \geq a$; therefore, $p(x) = 0$ for $x \geq a$. Assertions (i) and (ii) immediately follow from these remarks.

Next, we see from the representation $\sigma(t, x) = (1/\pi)(a - t - x)/\sqrt{t(a - t)}$ that for $x \in (0, a)$ the function $\sigma(t, x)$ is positive for $t \in [0, a - x)$ and negative in $(a - x, a]$. Hence,

$$\begin{aligned} p(x, a) &= \frac{1}{\pi} \int_0^{a-x} \sigma(t, x) dt = \frac{1}{\pi} \int_0^{a-x} \sqrt{\frac{a-t}{t}} dt - \frac{x}{\pi} \int_0^{a-x} \frac{dt}{\sqrt{t(a-t)}} \\ &= \frac{a}{\pi} \int_0^b \sqrt{\frac{1-t}{t}} dt - \frac{x}{\pi} \int_0^b \frac{dt}{\sqrt{t(1-t)}}, \end{aligned} \quad (2.16)$$

where

$$b := 1 - \frac{x}{a}.$$

Using the identities

$$\int_0^b \sqrt{\frac{1-t}{t}} dt = \sin^{-1} \sqrt{b} + \sqrt{b(1-b)}$$

and

$$\int_0^b \frac{dt}{\sqrt{t(1-t)}} = 2 \sin^{-1} \sqrt{b},$$

we obtain assertion (iii) from (??).

The first equality in (iv) may be obtained by differentiation of (iii); the second inequality then follows from the first and (i). ■

We now investigate the set of values of $a > 0$ satisfying the condition in the assertion of Lemma ???. For given $\alpha, \beta \geq 0, \alpha + \beta > 0$ we set

$$A := A(\alpha, \beta) := \{a : \exists x \in \mathbf{R} \text{ with } p(x, a) \leq \beta, n(x, a) \leq \alpha\}. \quad (2.17)$$

Lemma 2.3 *For any $\alpha, \beta \geq 0, \alpha + \beta > 0$ and $a > 0$ there exists a unique root $\bar{x} = \bar{x}(a; \alpha, \beta)$ of the equation*

$$\alpha p(x, a) = \beta n(x, a) \quad (2.18)$$

in $[0, a]$. Furthermore,

$$A = \{a > 0 : p(\bar{x}, a) \leq \beta\} \quad \text{for } \beta > 0, \quad (2.19a)$$

$$A = \{a > 0 : n(\bar{x}, a) \leq \alpha\} \quad \text{for } \alpha > 0, \quad (2.19b)$$

where A is defined in (??).

Proof. Consider first the case $\alpha, \beta > 0$. It follows by Lemma ?? that for fixed $a > 0$ the function $p(x, a)$ decreases on $(-\infty, a)$ from $+\infty$ to 0 and

$p(x, a) = 0$ for $x \geq a$. Also, the function $n(x, a)$ increases from 0 to $+\infty$ on $(0, \infty)$ and $n(x, a) = 0$ for $x \leq 0$. The same is true for $(\beta/\alpha)n(x, a)$ and therefore there exists a unique root \bar{x} of the equation

$$p(x, a) = \frac{\beta}{\alpha}n(x, a), \quad x \in (0, a), \quad (2.20)$$

which is equivalent to (??) for $\alpha, \beta > 0$. It also follows from the behavior of $n(x, a), p(x, a)$ that

$$p(\bar{x}, a) = \frac{\beta}{\alpha}n(\bar{x}, a) = \min_{x \in \mathbf{R}} \max\{p(x, a), \frac{\beta}{\alpha}n(x, a)\}.$$

On the other hand, the definition (??) may be written as

$$A = \{a : \min_{x \in \mathbf{R}} \max\{p(x, a), \frac{\beta}{\alpha}n(x, a)\} \leq \beta\}$$

provided $\alpha, \beta > 0$. Assertions (??) and (??) follow by these remarks.

It remains to notice that for $\alpha = 0, \beta > 0$ the representation (??) holds with $\bar{x} = 0$, which is the unique root of (??) in $[0, a]$ for $\alpha = 0$. Similarly, for $\alpha > 0, \beta = 0$ the representation (??) is true with $\bar{x} = a$, which is the unique root of (??) in $[0, a]$ for $\alpha > 0, \beta = 0$. ■

We define the function $G(y)$ for $y \in [0, 1]$ by

$$\pi G(y) := (1 - 2y) \sin^{-1} \sqrt{1 - y} + \sqrt{y(1 - y)}, \quad y \in [0, 1]. \quad (2.21)$$

Lemma 2.4 *The equation*

$$(\alpha - \beta)G(y) = \beta \left(y - \frac{1}{2} \right) \quad (2.22)$$

has a unique root $\bar{y} = \bar{y}(\alpha, \beta)$ in $[0, 1]$. Furthermore, with \bar{x} as in Lemma ??

$$\bar{x}(a; \alpha, \beta) = a\bar{y}(\alpha, \beta), \quad (2.23)$$

$$p(x, a) = aG(x/a), \quad a > 0, x \in [0, a]. \quad (2.24)$$

Proof. For $\alpha = \beta$ the unique root of (??) is $\bar{y} = 1/2$. For $\alpha \neq \beta$ we can rewrite (??) as

$$G(y) = \left(y - \frac{1}{2} \right) \frac{\beta}{\alpha - \beta}. \quad (2.25)$$

The range of $\kappa := \beta/(\alpha - \beta)$ for $\alpha, \beta \geq 0$ is $(-\infty, -1] \cup [0, \infty)$. We also note that

$$G'(y) = -\frac{2}{\pi} \sin^{-1} \sqrt{1-y} \in [-1, 0], \quad \text{for } y \in [0, 1].$$

Now, if $\kappa \in (-\infty, -1]$, then the function

$$G_1(y) := G(y) - \kappa \left(y - \frac{1}{2} \right)$$

is increasing in $[0, 1]$ since $G'_1(y) = G'(y) - \kappa \geq 0$. We have also $G_1(0) = 1/2 + \kappa/2 \leq 0$ and $G_1(1/2) = 1/2\pi > 0$. Thus (??) has a unique root \bar{y} for $\beta/(\alpha - \beta) \in (-\infty, -1]$. Next, if $\kappa \in [0, \infty)$, then $G'_1(y) = G'(y) - \kappa \leq 0$

for $y \in [0, 1]$ and so $G_1(y)$ decreases on $[0, 1]$. Since $G_1(1/2) = 1/2\pi$ and $G_1(1) = -\kappa/2 \leq 0$, (??) has in $[0, 1]$ a unique root \bar{y} which actually belongs to $[1/2, 1]$.

The equality (??) follows by (??) and (iii) of Lemma ?? . It remains to compare equations (??) and (??). Using (i) of Lemma ?? we may rewrite (??) as

$$(-\beta + \alpha)p(x, a) = \beta \left(x - \frac{a}{2} \right)$$

or, using (??), as

$$(-\beta + \alpha)G\left(\frac{x}{a}\right) = \beta \left(\frac{x}{a} - \frac{1}{2} \right).$$

The last equation coincides with (??) for $y = x/a$. Since both have a unique solution we obtain the assertion (??). ■

Lemma 2.5 *For any $\alpha, \beta \geq 0$, $\alpha + \beta > 0$ we have (cf. (??))*

$$A = [0, \bar{a}], \tag{2.26}$$

where $\bar{a} := \bar{a}(\alpha, \beta)$ is defined by

$$\bar{a} = \frac{\beta}{G(\bar{y})} = \frac{\alpha - \beta}{\bar{y} - 1/2} \quad \text{for } \alpha \neq \beta, \tag{2.27}$$

$$\bar{a} = 2\pi\beta \quad \text{for } \alpha = \beta. \tag{2.28}$$

Proof. Let $\alpha = \beta$. Then (??) has the unique root $\bar{y} = 1/2$ and $G(\bar{y}) = G(1/2) = 1/2\pi$ by (??). Now, it follows by (??) and (??) that $p(\bar{x}, a) = a/2\pi$ and so (??) may be written as $A = \{a > 0 : a/2\pi \leq \beta\}$. Assertion (??) follows.

Suppose $\alpha \neq \beta$ and $\beta > 0$. It follows by (??) and (??) that (??) may be written as $A = \{a : aG(\bar{y}) \leq \beta\}$. Then (??) follows from the second equality in (??) and the fact that $\alpha > \beta$ implies that $\bar{y} > 1/2$ (cf. the proof of Lemma ??).

In case $\beta = 0$ we rewrite (??) using (i) of Lemma ?? as

$$A = \left\{ a > 0 : p(\bar{x}, a) + \bar{x} - \frac{a}{2} \leq \alpha \right\}. \quad (2.29)$$

On the other hand, we have by (??) and (??) that

$$\begin{aligned} p(\bar{x}, a) + \bar{x} - \frac{a}{2} &= a \left\{ G(\bar{y}) + \left(\bar{y} - \frac{1}{2}\right) \right\} = a \left(\frac{\beta}{\alpha - \beta} + 1 \right) \left(\bar{y} - \frac{1}{2} \right) \\ &= a \cdot \frac{\alpha}{\alpha - \beta} \left(\bar{y} - \frac{1}{2} \right). \end{aligned}$$

Therefore (??) is equivalent to (??). ■

Proof of Theorem 1.3. Using the notation of (??), the assertion of Lemma ?? may be written as follows. If $a < a^*$, then $a \in A$. In view of

Lemma ?? this means that

$$(a < a^*) \implies (a \leq \bar{a})$$

and therefore we have $a^* \leq \bar{a}$.

It remains to notice from (??) and (1.8) that $\hat{y} = 1 - \bar{y}$, and so from (??) and (1.7) we have $\hat{a} = \bar{a}$. ■

3 Proofs of Theorems 1.2 and 1.4

For a fixed $b \in (0, 1)$ we let

$$\tilde{\sigma}_0(t) := \frac{1}{\pi} \frac{1}{\sqrt{(t-b)(1-t)}}, \quad t \in [b, 1], \quad (3.1)$$

$$\tilde{\sigma}_1(t) := \sqrt{b} \frac{\tilde{\sigma}_0(t)}{t}, \quad t \in [b, 1], \quad (3.2)$$

$$\tilde{\sigma}(t, x) := \tilde{\sigma}_1(t) - x\tilde{\sigma}_0(t), \quad t \in [b, 1], \quad x \in \mathbf{R}. \quad (3.3)$$

We note that $\tilde{\sigma}_0 dt$ is the equilibrium distribution for $[b, 1]$ and $\tilde{\sigma}_1 dt$ is the balayage of the unit mass at $x = 0$ to $[b, 1]$. Thus we have

$$V(x, \tilde{\sigma}_0 dt) = \log \frac{4}{1-b}, \quad x \in [b, 1], \quad (3.4)$$

$$V(x, \tilde{\sigma}_1 dt) = \log \frac{1}{x} + \text{const}, \quad x \in [b, 1]. \quad (3.5)$$

For the measure $\tilde{\sigma}(t, x)dt$ ($x \in \mathbf{R}$ is fixed) we consider its Jordan decomposition $\tilde{\sigma}dt = \tilde{\sigma}^+dt - \tilde{\sigma}^-dt$ and put

$$p(x) := p(x, b) := \int_{\Delta} \tilde{\sigma}^+(t, x)dt, \quad (3.6)$$

$$n(x) := n(x, b) := \int_{\Delta} \tilde{\sigma}^-(t, x)dt, \quad (3.7)$$

where $\Delta := [b, 1]$.

Following the scheme of proof of Lemma ?? we obtain the following.

Lemma 3.1 *Let b^* be defined as in (??). If $1 > b > b^*$, then there exists an $x \in \mathbf{R}$ such that*

$$p(x, b) \leq \frac{\beta}{\theta} \quad \text{and} \quad n(x, b) \leq \frac{\alpha}{\theta}.$$

Lemma 3.2 *The functions $p(x, b)$ and $n(x, b)$ defined in (??) and (??) satisfy the following properties:*

- (i) $p(x, b) - n(x, b) = 1 - x, \quad x \in \mathbf{R}, b \in (0, 1);$
- (ii) $p(x, b) = 1 - x$ and $n(x, b) = 0$ for $x \leq \sqrt{b}$,
 $p(x, b) = 0$ and $n(x, b) = x - 1$ for $x \geq 1/\sqrt{b}$;
- (iii) $p(x, b) = \int_b^{\sqrt{b}/x} \tilde{\sigma}(t, x)dt, \quad \sqrt{b} \leq x \leq 1/\sqrt{b};$

$$(iv) \quad \frac{\partial}{\partial x} p(x, b) = -\frac{2}{\pi} \sin^{-1} \sqrt{\frac{\sqrt{b} (1 - x\sqrt{b})}{x(1-b)}}, \quad \sqrt{b} < x \leq 1/\sqrt{b};$$

$$(v) \quad p(x, b) = \frac{2}{\pi} \int_x^{1/\sqrt{b}} \sin^{-1} \sqrt{\frac{\sqrt{b} (1 - t\sqrt{b})}{t(1-b)}} dt, \quad \sqrt{b} \leq x \leq 1/\sqrt{b};$$

$$(vi) \quad p(x, b) = \frac{2}{\pi} \sin^{-1} \sqrt{\frac{1 - x\sqrt{b}}{1-b}} - x \frac{2}{\pi} \sin^{-1} \sqrt{\frac{\sqrt{b} (1 - x\sqrt{b})}{x(1-b)}}, \quad \sqrt{b} \leq x \leq 1/\sqrt{b};$$

$$(vii) \quad p(x, b) = \frac{2}{\pi} \tan^{-1} \sqrt{\frac{1 - x\sqrt{b}}{\sqrt{b} (x - \sqrt{b})}} - x \frac{2}{\pi} \tan^{-1} \sqrt{\frac{\sqrt{b} (1 - x\sqrt{b})}{x - \sqrt{b}}}, \quad \sqrt{b} \leq x \leq 1/\sqrt{b};$$

$$(viii) \quad \frac{\partial}{\partial b} p(x, b) = -\frac{1}{\pi} \sqrt{\frac{(x - \sqrt{b})(1 - x\sqrt{b})}{b^{3/4}(1-b)}}, \quad \sqrt{b} \leq x \leq 1/\sqrt{b};$$

$$(ix) \quad p(x, b) = 1 - \frac{1}{\pi} \int_0^b \frac{\sqrt{(x - \sqrt{t})(1 - x\sqrt{t})}}{t^{3/4}(1-t)} dt, \quad \sqrt{b} \leq x \leq 1/\sqrt{b}.$$

Proof.

(i) With $\Delta = [b, 1]$, we have

$$p(x, b) - n(x, b) = \int_{\Delta} \tilde{\sigma}(t, x) dt = \int_{\Delta} \tilde{\sigma}_1(t) dt - x \int_{\Delta} \tilde{\sigma}_0(t) dt = 1 - x.$$

(ii) The function $\tilde{\sigma}(t, x) = (\sqrt{b}/t - x) \tilde{\sigma}_0(t)$ satisfies the inequalities $\tilde{\sigma}(t, x) \geq 0$ for $t \in \Delta$ if $x \leq \sqrt{b}$ and $\tilde{\sigma}(t, x) \leq 0$ for $t \in \Delta$ if $x \geq 1/\sqrt{b}$. Hence (ii)

follows from (i).

(iii) This property follows immediately from the definition of p .

(iv) We use (iii) to take derivative with respect to x . Since $\tilde{\sigma}(\sqrt{b}/x, x) = 0$,

we obtain

$$\begin{aligned} \frac{\partial p}{\partial x}(x, b) &= - \int_b^{\sqrt{b}/x} \tilde{\sigma}_0(t) dt = -\frac{1}{\pi} \int_b^{\sqrt{b}/x} \frac{dt}{\sqrt{(t-b)(1-t)}} \\ &= -\frac{1}{\pi} \int_0^\lambda \frac{d\tau}{\sqrt{\tau(1-\tau)}} = -\frac{2}{\pi} \sin^{-1} \sqrt{\lambda}, \end{aligned}$$

where

$$\lambda := \lambda(x, b) = \frac{\sqrt{b}/x - b}{1-b} = \frac{\sqrt{b}1 - x\sqrt{b}}{x1 - b} \quad (3.8)$$

(we use this notation hereafter).

(v) Since $p(1/\sqrt{b}, b) = 0$, (v) follows from (iv).

(vi) We integrate by parts in (v) to obtain

$$\begin{aligned} \frac{\pi}{2} p(x, b) &= \int_x^{1/\sqrt{b}} \sin^{-1} \sqrt{\lambda} dt = t \sin^{-1} \sqrt{\lambda} \Big|_x^{1/\sqrt{b}} - \int_x^{1/\sqrt{b}} t \frac{d}{dt} (\sin^{-1} \sqrt{\lambda}) dt \\ &= -x \sin^{-1} \sqrt{\lambda(x, b)} - \frac{1}{2} \int_x^{1/\sqrt{b}} \frac{t(\partial\lambda/\partial t)(t, b)}{\sqrt{\lambda(1-\lambda)}} dt. \end{aligned}$$

For the integrand in the last term we have

$$\frac{t\partial\lambda/\partial t}{\sqrt{\lambda(1-\lambda)}} = \frac{-1}{\sqrt{(1/\sqrt{b}-t)(t-\sqrt{b})}},$$

and after the substitution $\tau = (1/\sqrt{b}-t)/(1/\sqrt{b}-\sqrt{b})$ in the inte-

gral we obtain

$$-\frac{1}{2} \int_x^{1/\sqrt{b}} \frac{t\partial\lambda/\partial t dt}{\sqrt{\lambda(1-\lambda)}} = \frac{1}{2} \int_x^{1/\sqrt{b}} \frac{dt}{\sqrt{(1/\sqrt{b}-t)(t-\sqrt{b})}}$$

$$= \frac{1}{2} \int_0^{\frac{1-x\sqrt{b}}{1-b}} \frac{d\tau}{\sqrt{\tau(1-\tau)}} = \sin^{-1} \sqrt{\frac{1-x\sqrt{b}}{1-b}}$$

and (vi) follows.

(vii) Since $\sin^{-1} \alpha = \tan^{-1} \frac{\alpha}{\sqrt{1-\alpha^2}}$, (vii) follows from (vi).

(viii) We use (v): $p(x, b) = (2/\pi) \int_x^{1/\sqrt{b}} \sin^{-1} \sqrt{\lambda(t, b)} dt$ to take the derivative with respect to b . We have

$$\frac{\partial}{\partial b} p(x, b) = \frac{2}{\pi} \int_x^{1/\sqrt{b}} \frac{\partial}{\partial b} \left(\sin^{-1} \sqrt{\lambda(t, b)} \right) dt = \frac{1}{\pi} \int_x^{1/\sqrt{b}} \frac{(\partial \lambda / \partial b)(t, b) dt}{\sqrt{\lambda(t, b)(1-\lambda(t, b))}},$$

since $\lambda(t, b)$ equals zero for $t = 1/\sqrt{b}$ (cf. (??)). The integrand in the last term is

$$\begin{aligned} \frac{\partial \lambda / \partial b}{\sqrt{\lambda(1-\lambda)}} &= \frac{(1/\sqrt{b} + \sqrt{b}) - 2t}{2t(1-b)^2} / \sqrt{\frac{\sqrt{b}(1-t\sqrt{b})(t-\sqrt{b})}{t^2(1-b)^2}} \\ &= \frac{1}{2\sqrt{b}(1-b)} \frac{\gamma - 2t}{\sqrt{\gamma t - t^2 - 1}}, \end{aligned}$$

where

$$\gamma := \frac{1}{\sqrt{b}} + \sqrt{b}.$$

Hence, we have

$$\frac{\partial}{\partial b} p(x, b) = \frac{1}{2\pi\sqrt{b}(1-b)} \int_x^{1/\sqrt{b}} \frac{\gamma - 2t}{\sqrt{\gamma t - t^2 - 1}} dt$$

$$= -\frac{\sqrt{\gamma x - x^2 - 1}}{\pi\sqrt{b}(1-b)}$$

(since $\gamma t - t^2 - 1 = 0$ for $t = 1/\sqrt{b}$). The representation (viii) now follows.

(ix) We have $p(x, 0) = 1$ from (vi). Now if $\sqrt{b} \leq x \leq 1/\sqrt{b}$, then

$\sqrt{t} \leq x \leq 1/\sqrt{t}$ for $t \in [0, b]$ and we can integrate (viii) with respect to t instead of b over $[0, b]$. ■

Next we define

$$\begin{aligned} B &= B(\theta; \alpha, \beta) \\ &:= \left\{ b \in (0, 1) : \exists x \in \mathbf{R} \text{ with } p(x, b) \leq \frac{\beta}{\theta}, \quad n(x, b) \leq \frac{\alpha}{\theta} \right\} \end{aligned} \tag{3.9}$$

and we set

$$\bar{b} = \bar{b}(\theta, \alpha, \beta) := \inf B. \tag{3.10}$$

Lemma 3.3 (i) *If $\beta/\theta \geq 1$, then $\bar{b} = 0$.*

(ii) *If $\beta/\theta < 1$, then \bar{b} is the unique root of the equation*

$$p\left(1 - \frac{\beta}{\theta} + \frac{\alpha}{\theta}, b\right) = \frac{\beta}{\theta}$$

satisfying $\sqrt{b} \leq 1 - \beta/\theta + \alpha/\theta \leq 1/\sqrt{b}$.

Proof. We consider first the case $\alpha, \beta > 0$. Using the notation

$$h(b) := \min_{x \in \mathbf{R}} \max \left\{ p(x, b), \frac{\beta}{\alpha} n(x, b) \right\} \quad (3.11)$$

we can rewrite the definition (??) of B as

$$B = \left\{ b \in (0, 1) : h(b) \leq \frac{\beta}{\theta} \right\}. \quad (3.12)$$

It follows by (i) and (viii) of Lemma ?? that

$$\frac{\partial}{\partial b} p(x, b) = \frac{\partial}{\partial b} n(x, b) < 0 \quad \text{for } \sqrt{b} < x < \frac{1}{\sqrt{b}}; \quad (3.13)$$

hence both functions (of b) $p(x, b)$, $n(x, b)$ decrease for fixed x in the indicated domain.

On the other hand, it follows by (i), (ii) and (iv) of Lemma ?? that for fixed $b \in (0, 1)$ the function $p(x, b)$ decreases from $+\infty$ at $x = -\infty$ to 0 at $x = 1/\sqrt{b}$ and $n(x, b)$ increases from 0 at $x = \sqrt{b}$ to $+\infty$ at $x = +\infty$. This means that the equation

$$\alpha p(x, b) = \beta n(x, b) \quad (3.14)$$

has a unique root $x_1(b) \in (\sqrt{b}, 1/\sqrt{b})$ and

$$h(b) = p(x_1(b), b). \quad (3.15)$$

We note that $x_1(b)$ is a continuous function of b and we can conclude that for any fixed $b \in (0, 1)$, definition (??) may be written as

$$h(b) = \min_{x \in [x_1 - \epsilon, x_1 + \epsilon]} \max \left\{ p(x, b), \frac{\beta}{\alpha} n(x, b) \right\}, \quad (3.16)$$

where $x_1 = x_1(b)$. Moreover, this equality holds in some neighborhood of b with the same value of x_1 as defined by the original value of b . If $\epsilon > 0$ is small enough, then $(x_1 - \epsilon, x_1 + \epsilon) \times (b - \epsilon, b + \epsilon) \subset D := \{(x, b) : b \in (0, 1), \sqrt{b} < x < 1/\sqrt{b}\}$ since $(x_1(b), b) \in D$. Hence, it follows by (??) and (??) that $h(b)$ is decreasing in some neighborhood of $b \in (0, 1)$. Since $b \in (0, 1)$ is arbitrary, we conclude that $h(b)$ is decreasing on $(0, 1)$.

Next, we observe that $p(x, b) \rightarrow 1$ as $b \rightarrow 0$ uniformly over any interval $x \in [0, R]$ (cf. (ii) and (vi) of Lemma ??). Also, as $b \rightarrow 0$, $(\beta/\alpha)n(x, b) \rightarrow (\beta/\alpha)x$ (cf. (i) of Lemma ??), so that $x_1(b) \rightarrow \alpha/\beta$ and $h(b) = p(x_1(b), b) \rightarrow 1$. On the other hand, we have $0 \leq p(x_1(b), b) \leq p(\sqrt{b}, b) = 1 - \sqrt{b}$. Therefore, $h(b) = p(x_1(b), b) \rightarrow 0$ as $b \rightarrow 1$. Hence $h(b)$ decreases from 1 to 0 on $(0, 1)$.

Now, if $\beta/\theta \geq 1$, then $h(b) \leq \beta/\theta$ for any $b \in (0, 1)$ and assertion (i) of the lemma follows by (??). If $\beta/\theta < 1$, then from the properties of h and p described above, the value of \bar{b} is determined by the system of equations

$$\alpha p(x, b) = \beta n(x, b), \quad p(x, b) = \frac{\beta}{\theta}, \quad (3.17)$$

which has the unique solution $(x_1(\bar{b}), \bar{b})$. Using (i) of Lemma ?? the first equation may be written as $p(x) = (\beta/\alpha)\{p(x)+x-1\}$ or $p(x)=\beta(x-1)/(\alpha-\beta)$. If we substitute this expression for $p(x)$ in the second equation of (??) we see that the system (??) is equivalent to

$$\begin{cases} x = 1 - \beta/\theta + \alpha/\theta, \\ p(x, b) = \beta/\theta. \end{cases}$$

Hence assertion (ii) of the lemma follows.

Now suppose that $\alpha = 0$, $\beta > 0$. The requirement $n(x, b) \leq 0$ is included in the definition (??) of the set B , which implies that $x \leq \sqrt{b}$. The minimal value for $p(x, b)$ over $(-\infty, \sqrt{b}]$ is achieved at $x = \sqrt{b}$; hence (??) may be written in this case as

$$B = \left\{ b \in (0, 1) : p(\sqrt{b}, b) \leq \frac{\beta}{\theta} \right\}.$$

Since $p(\sqrt{b}, b) = 1 - \sqrt{b}$, this yields for $\bar{b} = \inf\{b \in B\}$,

$$\bar{b} = 0 \text{ if } \beta/\theta \geq 1, \quad \bar{b} = \left(1 - \frac{\beta}{\theta}\right)^2 \text{ if } \beta/\theta < 1. \quad (3.18)$$

For $\beta/\theta < 1$, it is clear that $\bar{b} = (1 - \beta/\theta)^2$ is a root of the equation $p(1 - \beta/\theta, b) = \beta/\theta$. It remains to show that this root is unique; i.e.

$$p(x, b) \neq 1 - x \quad \text{for } \sqrt{b} < x \leq 1/\sqrt{b}. \quad (3.19)$$

It follows from (iv) of Lemma ?? that

$$\frac{\partial}{\partial x} p(x, b) > -1 \quad \text{for } \sqrt{b} < x \leq 1/\sqrt{b}.$$

Therefore,

$$\begin{aligned} p(x, b) &= p(\sqrt{b}, b) + \int_{\sqrt{b}}^x \frac{\partial}{\partial t} p(t, b) dt \\ &> 1 - \sqrt{b} - (x - \sqrt{b}) = 1 - x, \quad \text{for } \sqrt{b} < x \leq 1/\sqrt{b}, \end{aligned}$$

which yields (?). Thus assertions (i) and (ii) of the lemma hold when $\alpha = 0$.

Finally, for $\alpha > 0, \beta = 0$, we have from definition (??) that for $b \in B$, there exists an x such that $p(x, b) \leq 0$, and so $x \geq 1/\sqrt{b}$. The minimal value for $n(x, b)$ over $x \geq 1/\sqrt{b}$ is achieved only at $x = 1/\sqrt{b}$ and thus $B = \{b \in (0, 1) : n(1/\sqrt{b}, b) \leq \alpha/\theta\}$ or

$$\bar{b} = \inf B = \frac{1}{(1 + \alpha/\theta)^2}. \quad (3.20)$$

Clearly $\bar{b} = (1 + \alpha/\theta)^{-2}$ is a root of $p(1 + \alpha/\theta, b) = 0$ satisfying $\sqrt{\bar{b}} \leq (1 + \alpha/\theta) \leq 1/\sqrt{\bar{b}}$. Moreover, according to (ii) and (v) of Lemma ??, \bar{b} is the only root of $p(1 + \alpha/\theta, b) = 0$ satisfying $\sqrt{\bar{b}} \leq 1 + \alpha/\theta \leq 1/\sqrt{\bar{b}}$. This completes the proof of Lemma ??. ■

Proof of Theorem ??. The assertion of Lemma ?? combined with definition of (??) may now be written as $(1 > b > b^*) \Rightarrow (b \in B) \Rightarrow (b \geq \bar{b})$.

This implies that

$$b^* \geq \bar{b}.$$

Using (ix) of Lemma ?? the equation for \bar{b} in (ii) of Lemma ?? may be written as

$$\frac{1}{\pi} \int_0^{\bar{b}} \frac{\sqrt{(\xi - \sqrt{t})(1 - \xi\sqrt{t})}}{t^{3/4}(1-t)} dt = 1 - \frac{\beta}{\theta} \quad (3.21)$$

with $\xi = 1 - \beta/\theta + \alpha/\theta$. Thus $\bar{b} = \hat{b}$ and the proof of Theorem ?? is complete. ■

Proof of Theorem ??. In the case $\alpha = \beta$ we have $\xi = 1 - \beta/\theta + \alpha/\theta = 1$ and the integral on the left-hand side of (??) becomes

$$\begin{aligned} \frac{1}{\pi} \int_0^{\bar{b}} \frac{1 - \sqrt{t}}{t^{3/4}(1-t)} dt &= \frac{1}{\pi} \int_0^{\bar{b}} \frac{dt}{t^{3/4}(1 + \sqrt{t})} = \frac{4}{\pi} \int_0^{\bar{b}} \frac{d(t^{1/4})}{1 + t^{1/2}} \\ &= \frac{4}{\pi} \tan^{-1}(\bar{b}^{1/4}). \end{aligned}$$

Hence, equation (??) has the solution

$$\bar{b} = \tan^4 \left(\frac{\pi}{4} \left(1 - \frac{\beta}{\theta} \right) \right).$$

Theorem ?? now follows from Theorem ?? (with $\alpha = \beta = 1$). ■

4 Appendix

Here we prove the identity (2.2):

$$u(x) := \frac{1}{\pi} \int_0^a \log \frac{1}{|x-t|} \sqrt{\frac{a-t}{t}} dt = -x + c, \quad x \in [0, a], \quad (\text{A.1})$$

where c is a constant.

The integral in (??) defines a function $u(x)$ continuous in the whole plane

C. In the upper half-plane, we have

$$u(x) = \operatorname{Re} U(x), \quad \operatorname{Im} x > 0, \quad (\text{A.2})$$

where

$$U(z) := -\frac{1}{\pi} \int_0^a \log(z-t) \sqrt{\frac{a-t}{t}} dt, \quad \operatorname{Im} z > 0, \quad (\text{A.3})$$

and the branch of log is determined by the normalization $0 < \arg(z-t) < \pi$,

$t \in [0, a], \operatorname{Im} z > 0$. The derivative

$$U'(z) = \frac{1}{\pi} \int_0^a \sqrt{\frac{a-t}{t}} \frac{dt}{t-z}, \quad z \in D := \overline{\mathbb{C}} \setminus [0, a], \quad (\text{A.4})$$

is a single-valued analytic function in D . Let

$$f(z) := \sqrt{\frac{z-a}{z}}, \quad z \in D, \quad f(\infty) := 1,$$

and denote by $f^+(x), f^-(x)$, $x \in [0, a]$, the boundary values of $f(z)$ from the upper and lower half-planes, respectively. Then we have

$$f^\pm(x) = \pm i \sqrt{\frac{a-x}{x}}, \quad (\text{A.5})$$

Using (??) we can rewrite (??) as

$$U'(z) = \frac{1}{2\pi i} \int_{\partial D} \frac{f(t)}{t-z} dt, \quad z \in D, \quad (\text{A.6})$$

where ∂D is the boundary of D with positive orientation with respect to D .

The integral in (??) is the Cauchy integral for $f(z)$ in D and therefore

$$U'(z) = f(z) - f(\infty) = \sqrt{\frac{z-a}{z}} - 1,$$

so that

$$U(z) = \int_0^z \sqrt{\frac{\zeta-a}{\zeta}} d\zeta - z + \text{const.}, \quad \text{Im } z > 0.$$

Now (??) follows from the last representation and (??).

Finally we remark that (??) is well-known (cf. [Ts]) since $\sigma_0 dt$ is the equilibrium distribution for the interval $[0, a]$, which has logarithmic capacity $a/4$.

References

- [B] P. Borwein, *The size of $\{x : r'_n/r_n \geq 1\}$ and lower bounds for $\|e^{-x} - r_n\|$* . J. Approx. Theory. **36** (1982) 73–80.
- [BC] P. Borwein and W. Chen, *Incomplete rationals in the complex plane*. Constr. Approx., to appear.
- [BS] P. Borwein and E. B. Saff, personal communication.
- [G] M. v. Golitschek, *Approximation by incomplete polynomials*, Journ. Approx. Theory, **28** (1980), 155-160.
- [GR] A. A. Gonchar and E. A. Rakhmanov, *Equilibrium distributions and degree of rational approximation of analytic functions*, Math. USSR Sbornik, **62** (1989), 305-348.
- [La] N. S. Landkof, *Foundations of Modern Potential Theory*, Springer-Verlag, New York, N.Y., (1972).
- [Lo] G. G. Lorentz, *Approximation by incomplete polynomials*, In: Padé and Rational Approximation: Theory and Applications (Eds. E. B. Saff and R. S. Varga), Academic Press, New York (1977), 289-302.

- [LR] G. Lopez and E. A. Rakhmanov, *Rational approximations, orthogonal polynomials and equilibrium distributions*, In: Orthogonal Polynomials and their Applications (Proceedings, Segovia, 1986), Lecture Notes in Math, **1329**, Springer-Verlag, Heidelberg (1988), 125-157.
- [LS] D. S. Lubinsky and E. B. Saff, *Uniform and mean approximation by certain weighted polynomials, with applications*, Constr. Approx., **4** (1988), 21-64.
- [ST] E. B. Saff and V. Totik, *Logarithmic Potentials with External Fields*, Springer-Verlag (to appear).
- [SV] E. B. Saff and R. S. Varga, *Uniform approximation by incomplete polynomials*, Internat. J. Math. Soc., **1** (1978), 407-420.
- [To] V. Totik, *Weighted Approximation with Varying Weight*, Lecture Notes in Math., **1569**, Springer-Verlag, Heidelberg (1994).
- [Ts] M. Tsuji, *Potential Theory in Modern Function Theory*, Chelsea Pub., New York (1959).

P. Borwein

Department of Mathematics

Simon Fraser University

Burnaby, British Columbia

Canada V5A 1S6

email: pborwein@cecm.sfu.ca

E. B. Saff

Institute for Constructive Mathematics

Department of Mathematics

University of South Florida

Tampa, Fl 33620

U.S.A.

email: esaff@math.usf.edu

E. A. Rakhmanov

Steklov Institute

42 Vavilova St

Moscow

Russia

email: rakhman@mph.mian.su

email: rakhmano@math.usf.edu