

ON CYCLOTOMIC POLYNOMIALS WITH ± 1 COEFFICIENTS

PETER BORWEIN AND KWOK-KWONG STEPHEN CHOI

Abstract.

We characterize all cyclotomic polynomials of even degree with coefficients restricted to the set $\{+1, -1\}$. In this context a cyclotomic polynomial is any monic polynomial with integer coefficients and all roots of modulus 1. *Inter alia* we characterize all cyclotomic polynomials with odd coefficients.

The characterization is as follows. A polynomial $P(x)$ with coefficients ± 1 of even degree $N - 1$ is cyclotomic if and only if

$$P(x) = \pm \Phi_{p_1}(\pm x) \Phi_{p_2}(\pm x^{p_1}) \cdots \Phi_{p_r}(\pm x^{p_1 p_2 \cdots p_{r-1}}),$$

where $N = p_1 p_2 \cdots p_r$ and the p_i are primes, not necessarily distinct. Here $\Phi_p(x) := \frac{x^p - 1}{x - 1}$ is the p th cyclotomic polynomial.

We conjecture that this characterization also holds for polynomials of odd degree with ± 1 coefficients. This conjecture is based on substantial computation plus a number of special cases.

Central to this paper is a careful analysis of the effect of Graeffe's root squaring algorithm on cyclotomic polynomials.

Keywords: Cyclotomic polynomials, Mahler measure, Littlewood polynomials
AMS (1991) subject classification: Primary 11R09, Secondary 11Y99

Research of P. Borwein is supported, in part, by NSERC of Canada. K.K. Choi is a Pacific Institute of Mathematics Postdoctoral Fellow and the Institute's support is gratefully acknowledged.

1. INTRODUCTION.

We are interested in studying polynomials with coefficients restricted to the set $\{+1, -1\}$. This particular set of polynomials has drawn much attention and there are a number of difficult old questions concerning it. Littlewood raised a number of these questions and so we call these polynomials *Littlewood polynomials* and denote them by \mathcal{L} [6]. A Littlewood polynomial of degree n has L_2 norm on the unit circle equal to $\sqrt{n+1}$. Many of the questions raised concern comparing the behavior of these polynomials in other norms to the L_2 norm. One of the older and more intriguing of these asks whether such polynomials can be “flat”. Specifically, do there exist two positive constants C_1 and C_2 so that for each n there is Littlewood polynomial of degree n with

$$C_1\sqrt{n+1} < |p(z)| < C_2\sqrt{n+1}$$

for each z of modulus 1. This problem which has been open for more than forty years is discussed in [1] where there is an extensive bibliography. The upper bound is satisfied by the so called Rudin–Shapiro polynomials. It is still open as to whether there is a sequence satisfying just the lower bound (this problem has been called one of the “very hardest problems in combinatorial optimization”).

The size of the L_p norm of Littlewood polynomials has been studied from a number of points of view. The problem of minimizing the L_4 norm (or equivalently of maximizing the so called “merit factor”) has also attracted a lot of attention.

In particular can Littlewood polynomials of degree n have L_4 norm asymptotically close to $\sqrt{n+1}$. This too is still open and is discussed in [1].

Mahler [7] raised the question of maximizing the Mahler measure of Littlewood polynomials. The Mahler measure is just the L_0 norm on the circle and one would expect this to be closely related to the minimization problem for the L_4 norm above. Of course the minimum possible Mahler measure for a Littlewood polynomial is 1 and this is achieved by any cyclotomic polynomial. In this paper a cyclotomic polynomial is any monic polynomial with integer coefficients and all roots of modulus 1. While $\Phi_n(x)$ denotes *the* n th irreducible cyclotomic polynomial (the minimum polynomial of a primitive n th root of unity).

This paper addresses the question of characterizing the cyclotomic Littlewood polynomials of even degree. Specifically we show that a polynomial $P(x)$ with coefficients ± 1 of even degree $N-1$ is cyclotomic if and only if

$$P(x) = \pm \Phi_{p_1}(\pm x) \Phi_{p_2}(\pm x^{p_1}) \cdots \Phi_{p_r}(\pm x^{p_1 p_2 \cdots p_{r-1}}),$$

where $N = p_1 p_2 \cdots p_r$ and the p_i are primes (not necessarily distinct). The “if” part is obvious since $\Phi_{p_i}(x)$ has coefficients ± 1 .

We also give an explicit formula for the number of such polynomials.

This analysis is based on a careful treatment of Graeffe’s root squaring algorithm. It transpires that all cyclotomic Littlewood polynomials of a fixed degree have the same fixed point on iterating Graeffe’s root squaring algorithm. This allows us to also characterize all cyclotomic polynomials with odd coefficients.

Substantial computations, as well as a number of special cases, lead us to conjecture that the above characterization of cyclotomic Littlewood polynomials of even degree also holds for odd degree. One of the cases we can handle is when N is a power of 2.

It is worth commenting on the experimental aspects of this paper. (As is perhaps usual, much of this is carefully erased in the final exposition). It is really the observation that the cyclotomic Littlewood polynomials can be explicitly constructed essentially by inverting Graeffe's root squaring algorithm that is critical. This allows for computation over all cyclotomic Littlewoods up to degree several hundred (with exhaustive search failing far earlier). A construction which is of interest in itself. Indeed it was these calculations that allowed for the conjectures of the paper and suggested the route to some of the results.

The paper is organized as follows. Section 2 examines cyclotomic polynomials with odd coefficients. Section 3 looks at cyclotomic Littlewood polynomials with a complete analysis of the even degree case. The last section presents some numerical evidence and other evidence to support the conjecture in the odd case behaves like the even case.

2. CYCLOTOMIC POLYNOMIALS WITH ODD COEFFICIENTS.

In this section, we discuss the factorization of cyclotomic polynomials with odd coefficients as a product of irreducible cyclotomic polynomials. To do this, we first consider the factorization over $\mathbb{Z}_p[x]$ where p is a prime number. The most useful case is $p = 2$ because every Littlewood polynomial reduces to the Dirichlet kernel $1 + x + \cdots + x^{N-1}$ in $\mathbb{Z}_2[x]$. In $\mathbb{Z}_p[x]$, $\Phi_n(x)$ is no longer irreducible in general but $\Phi_n(x)$ and $\Phi_m(x)$ are still relatively prime to each other. Here, as before, $\Phi_n(x)$ is the n th irreducible cyclotomic polynomial.

Lemma 2.1. *Suppose n and m are distinct positive integers relatively prime to p . Then $\Phi_n(x)$ and $\Phi_m(x)$ are relatively prime in $\mathbb{Z}_p[x]$.*

Proof. Suppose e and f are the smallest positive integers such that

$$p^e \equiv 1 \pmod{n} \quad \text{and} \quad p^f \equiv 1 \pmod{m}.$$

Let F_{p^k} be the field of order p^k . Then F_{p^e} contains exactly $\phi(n)$ elements of order n and over \mathbb{Z}_p , $\Phi_n(x)$ is a product of $\phi(n)/e$ irreducible factors of degree e and each irreducible factor is a minimal polynomial for an element in F_{p^e} of order n over \mathbb{Z}_p , see [5]. So $\Phi_n(x)$ and $\Phi_m(x)$ cannot have a common factor in $\mathbb{Z}_p[x]$ since their irreducible factors are minimal polynomials of different orders. This proves our lemma. \square

The following lemma tells which $\Phi_m(x)$ can possibly be factors of polynomials with odd coefficients.

Lemma 2.2. *Suppose $P(x)$ is a polynomial with odd coefficients of degree $N - 1$. If $\Phi_m(x)$ divides $P(x)$, then m divides $2N$.*

Proof. Since $\Phi_m(x)$ divides $P(x)$, so $\Phi_m(x)$ also divides $P(x)$ in $\mathbb{Z}_2[x]$. However, in $\mathbb{Z}_2[x]$, $P(x)$ equals to $1 + x + \cdots + x^{N-1}$ and can be factored as

$$(2.1) \quad P(x) = \Phi_1(x)^{-1} \prod_{d|M} \Phi_d^{2^t}(x),$$

where $N = 2^t M$, $t \geq 0$ and M is odd. In view of Lemma 2.1, $\Phi_{d_1}(x)$ and $\Phi_{d_2}(x)$ are relatively prime in $\mathbb{Z}_2[x]$ if d_1 and d_2 are distinct odd integers. So if m is odd, then $\Phi_m(x)$ is a factor in the right hand side of (2.1) and hence $m = d$ for some

$d|M$. On the other hand, if m is even and $m = 2^l m'$ where $l \geq 1$ and m' is odd, then

$$\begin{aligned}\Phi_m(x) &= \Phi_{2m'}(x^{2^{l-1}}) \\ &= \Phi_{m'}(x^{2^{l-1}}) \\ &= \Phi_{m'}(x)^{2^{l-1}}\end{aligned}$$

in $\mathbb{Z}_2[x]$. Thus in view of (2.1), we must have $m' = d$ for $d|M$ and $l \leq t + 1$. Hence in both cases, we have m divides $2N$. \square

In view of Lemma 2.2, every cyclotomic polynomial, $P(x)$, with odd coefficients of degree $N - 1$ can be written as

$$(2.2) \quad P(x) = \prod_{d|2N} \Phi_d^{e(d)}(x)$$

where $e(d)$ are non-negative integers.

For each prime p let T_p be the operator defined over all monic polynomials in $\mathbb{Z}[x]$ by

$$T_p[P(x)] := \prod_{i=1}^N (x - \alpha_i^p)$$

for every $P(x) = \prod_{i=1}^N (x - \alpha_i)$ in $\mathbb{Z}[x]$. By Newton's identities (see [3], p.5), $T_p[P(x)]$ is also a monic polynomial in $\mathbb{Z}[x]$. We extend T_p to be defined over the quotient of two monic polynomials in $\mathbb{Z}[x]$ by $T_p[(P/Q)(x)] := T_p[P(x)]/T_p[Q(x)]$. This operator obviously takes a polynomial to the polynomial whose roots are the p th powers of the roots of P . Also we let M_p be the natural projection from $\mathbb{Z}[x]$ onto $\mathbb{Z}_p[x]$. So,

$$M_p[P(x)] = P(x) \pmod{p}.$$

Lemma 2.3. *Let n be a positive integer relatively prime to p and $i \geq 2$. Then we have*

- (i) $T_p[\Phi_n(x)] = \Phi_n(x)$,
- (ii) $T_p[\Phi_{pn}(x)] = \Phi_n(x)^{p-1}$,
- (iii) $T_p[\Phi_{p^i n}(x)] = \Phi_{p^{i-1}n}(x)^p$.

Proof. (i) is trivial because if $(n, p) = 1$ then T_p just permutes the roots of $\Phi_n(x)$. To prove (ii) and (iii), we consider

$$\begin{aligned}T_p[P(x^p)] &= T_p\left[\prod_{j=1}^N (x^p - \alpha_j)\right] \\ &= T_p\left[\prod_{j=1}^N \prod_{l=1}^p (x - e^{\frac{2\pi i l}{p}} \alpha_j^{\frac{1}{p}})\right] \\ &= \prod_{j=1}^N \prod_{l=1}^p (x - \alpha_j) \\ &= P(x)^p.\end{aligned}$$

Thus (ii) and (iii) follow from (i), $\Phi_{pn}(x) = \frac{\Phi_n(x^p)}{\Phi_n(x)}$ and $\Phi_{p^i n}(x) = \Phi_{p^{i-1}n}(x^p)$ (see [8], §5.8). \square

When $P(x)$ is cyclotomic, the iterates $T_p^n[P(x)]$ converge in a finite number of steps to a fixed point of T_p and we define this to be the fixed point of $P(x)$ with respect to T_p .

Lemma 2.4. *If $P(x)$ is a monic cyclotomic polynomial in $\mathbb{Z}[x]$, then*

$$(2.3) \quad M_p[T_p[P(x)]] = M_p[P(x)],$$

in $\mathbb{Z}_p[x]$, where M_p is the above natural projection.

Proof. Since both T_p and M_p are multiplicative, it suffices to consider the primitive cyclotomic polynomials $\Phi_n(x)$. Let n be an integer relatively prime to p . Then (2.3) is true for $P(x) = \Phi_n(x)$ by (i) of Lemma 2.3. For $P(x) = \Phi_{pn}(x)$, we have

$$\begin{aligned} M_p[T_p[\Phi_{pn}(x)]] &= M_p[\Phi_n(x)^{p-1}] \\ &= M_p[\Phi_n(x)]^{p-1}. \end{aligned}$$

by (ii) of Lemma 2.3. However,

$$\begin{aligned} M_p[\Phi_{pn}(x)] &= \frac{M_p[\Phi_n(x^p)]}{M_p[\Phi_n(x)]} \\ &= \frac{M_p[\Phi_n](x^p)}{M_p[\Phi_n(x)]} \\ &= M_p[\Phi_n(x)]^{p-1}, \end{aligned}$$

in $\mathbb{Z}_p[x]$. This proves that (2.3) is also true for $P(x) = \Phi_{pn}(x)$. Finally, if $P(x) = \Phi_{p^i n}(x)$ then

$$\begin{aligned} M_p[T_p[\Phi_{p^i n}(x)]] &= M_p[\Phi_{p^{i-1} n}(x)^p] \\ &= M_p[\Phi_{p^{i-1} n}(x^p)] \\ &= M_p[\Phi_{p^i n}(x)] \end{aligned}$$

by (iii) of Lemma 2.3. This completes the proof of our lemma. \square

Lemma 2.4 shows that if $T_p[P(x)] = T_p[Q(x)]$ then $M_p[P(x)] = M_p[Q(x)]$. The next result shows that the converse is also true.

Theorem 2.5. *$P(x)$ and $Q(x)$ are monic cyclotomic polynomials in $\mathbb{Z}[x]$ and $M_p[P(x)] = M_p[Q(x)]$ in $\mathbb{Z}_p[x]$ if and only if both $P(x)$ and $Q(x)$ have the same fixed point with respect to iteration of T_p .*

Proof. Suppose

$$P(x) = \prod_{d \in \mathcal{D}} \Phi_d^{e(d)}(x) \Phi_{pd}^{e(pd)}(x) \cdots \Phi_{p^t d}^{e(p^t d)}(x)$$

and

$$Q(x) = \prod_{d \in \mathcal{D}} \Phi_d^{e(d)'}(x) \Phi_{pd}^{e(pd)'}(x) \cdots \Phi_{p^t d}^{e(p^t d)'}(x)$$

where $t, e(j), e(j)' \geq 0$ and \mathcal{D} is a set of positive integers relatively prime to p . Then using (i)-(iii) of Lemma 2.3, we have for $l \geq t$

$$(2.4) \quad T_p^l[P(x)] = \prod_{d \in \mathcal{D}} \Phi_d(x)^{f(d)} \quad \text{and} \quad T_p^l[Q(x)] = \prod_{d \in \mathcal{D}} \Phi_d(x)^{f(d)'}$$

where

$$f(d) = e(d) + (p-1) \sum_{j=1}^t p^{j-1} e(p^j d)$$

and

$$f(d)' = e(d)' + (p-1) \sum_{j=1}^t p^{j-1} e(p^j d)'.$$

From Lemma 2.4, we have

$$M_p[T_p^l[P(x)]] = M_p[P(x)] = M_p[Q(x)] = M_p[T_p^l[Q(x)]],$$

for any $l \geq t$. From this and (2.4),

$$\prod_{d \in \mathcal{D}} M_p[\Phi_d(x)]^{f(d)} = \prod_{d \in \mathcal{D}} M_p[\Phi_d(x)]^{f(d)'}$$

However, with Lemma 2.1, $M_p[\Phi_d(x)]$ and $M_p[\Phi_{d'}(x)]$ are relatively prime if $d \neq d'$. So we must have $f(d) = f(d)'$ for all $d \in \mathcal{D}$ and hence from (2.4), $P(x)$ and $Q(x)$ have the same fixed point with respect to T_p . \square

From Theorem 2.5, we can characterize the monic cyclotomic polynomials by their images in $\mathbb{Z}_p[x]$ under the projection M_p . They all have the same fixed point under T_p . In particular, when $p = 2$ we have

Corollary 2.6. *All monic cyclotomic polynomials with odd coefficients of the degree $N - 1$ have the same fixed point under iteration of T_2 . Specifically, if $N = 2^t M$ where $t \geq 0$ and $(2, M) = 1$ then the fixed point occurs at the $t + 1$ -th step of the iteration and equals*

$$(x^M - 1)^{2^t} (x - 1)^{-1}.$$

Proof. The first part follows directly from our Theorem 2.5 and the fact that

$$M_2[P(x)] = 1 + x + \cdots + x^{N-1}$$

in $\mathbb{Z}_2[x]$ if $P(x)$ is a monic polynomials with odd coefficients of degree $N - 1$. If $N = 2^t M$, then from (2.2),

$$P(x) = \prod_{d|M} \Phi_d^{e(d)}(x) \Phi_{2d}^{e(2d)}(x) \cdots \Phi_{2^{t+1}d}^{e(2^{t+1}d)}(x).$$

Over $\mathbb{Z}_2[x]$,

$$1 + x + \cdots + x^{N-1} = \Phi_1(x)^{-1} \prod_{d|M} \Phi_d^{2^t}(x),$$

so

$$(2.5) \quad f(d) = e(d) + \sum_{i=1}^{t+1} 2^{i-1} e(2^i d) = \begin{cases} 2^t & \text{for } d|M, d > 1, \\ 2^t - 1 & \text{for } d = 1. \end{cases}$$

Therefore, from (2.5) and Lemma 2.3,

$$T_2^{t+1}[P(x)] = \prod_{d|M} \Phi_d^{f(d)}(x) = \Phi_1(x)^{-1} \prod_{d|M} \Phi_d^{2^t}(x) = (x^M - 1)^{2^t} (x - 1)^{-1}.$$

\square

Corollary 2.6, when N is odd ($t = 0$) shows that $T_2[P(x)]$ equals to $1 + x + \dots + x^{N-1}$ for all cyclotomic polynomials with odd coefficients and from (2.2) and (2.5), we have the following characterization of cyclotomic polynomials with odd coefficients.

Corollary 2.7. *Let $N = 2^t M$ with $t \geq 0$ and $(2, M) = 1$. A polynomial, $P(x)$, with odd coefficients of degree $N - 1$ is cyclotomic if and only if*

$$P(x) = \prod_{d|M} \Phi_d^{e(d)}(x) \Phi_{2d}^{e(2d)}(x) \dots \Phi_{2^{t+1}d}^{e(2^{t+1}d)}(x),$$

and the $e(d)$ satisfy the condition (2.5).

Furthermore, if N is odd, then any polynomial, $P(x)$, with odd coefficients of even degree $N - 1$ is cyclotomic if and only if

$$P(x) = \prod_{d|N, d>1} \Phi_d^{e(d)}(\pm x)$$

where the $e(d)$ are non-negative integers.

Corollary 2.7 allows us to compute the number of cyclotomic polynomials with odd coefficients. Let $B(n)$ be the number of partitions of n into a sum of terms of the sequence $\{1, 1, 2, 4, 8, 16, \dots\}$. Then $B(n)$ has generating function

$$F(x) = (1 - x)^{-1} \prod_{k=0}^{\infty} (1 - x^{2^k})^{-1}.$$

It follows from (2.5) and Corollary 2.7 that

Corollary 2.8. *Let $N = 2^t M$ with $t \geq 0$ and $(2, M) = 1$. The number of cyclotomic polynomials with odd coefficients of degree $N - 1$ is*

$$(2.6) \quad C(N) = B(2^t)^{d(M)-1} \cdot B(2^t - 1)$$

where $d(M)$ denotes the number of divisors of M . Furthermore,

$$(2.7) \quad \log C(N) \sim \left(\frac{t^2}{2} \log 2\right)(d(M) - 1) + \frac{(\log(2^t - 1))^2}{\log 4}.$$

Proof. Formula (2.6) follows from (2.5) and Corollary 2.7. To prove (2.7), we use de Bruijn's asymptotic estimation for $B(n)$ in [4]:

$$B(n) \sim \exp((\log n)^2 / \log 4).$$

Now (2.7) follows from this and (2.6). \square

3. CYCLOTOMIC LITTLEWOOD POLYNOMIALS.

We now specialize the discussion to the case where the coefficients are all plus one or minus one.

One natural way to build up Littlewood polynomial of higher degree is as follows: if $P_1(x)$ and $P_2(x)$ are Littlewood polynomials and $P_1(x)$ is of degree $N - 1$ then $P_1(x)P_2(x^N)$ is a Littlewood polynomials of higher degree. In this section, we are going to show that this is also the only way to produce cyclotomic Littlewood polynomials, at least, for even degree.

To prove this, it is equivalent to show that the coefficients of $P(x)$ are "periodic" in the sense that if $P(x) = \sum_{n=0}^{N-1} a_n x^n$, then there is a "period" i such that

$a_{li+n} = a_{li}$ for all $1 \leq n \leq i-1$ and $0 \leq l \leq N/i-1$. This is our Theorem 3.3 below.

Suppose $P(x) = \sum_{n=0}^{N-1} a_n x^n$ is a cyclotomic polynomial in \mathcal{L} and let S_k be the sum of k -th power of all the roots of $P(x)$. Since $P(x)$ is cyclotomic, we have $x^{N-1}P(1/x) = \pm P(x)$. Thus it follows from Newton's identities that

$$(3.1) \quad S_k + a_1 S_{k-1} + \cdots + a_{k-1} S_1 + k a_k = 0$$

for $k \leq N-2$. We may further assume that $a_0 = a_1 = 1$ by replacing $P(x)$ by $-P(x)$ or $P(-x)$ if necessary. We now let

$$a_0 = a_1 = \cdots = a_{i-1} = 1 \quad \text{and} \quad a_i = -1,$$

for some integer $i \geq 2$. From (3.1), we have

$$S_1 = -a_1 = -1.$$

We claim that

$$(3.2) \quad S_1 = S_2 = \cdots = S_{i-1} = -1 \quad \text{and} \quad S_i = 2i - 1.$$

Suppose $S_1 = \cdots = S_j = -1$ for $j < i-1$. Then from (3.1) again,

$$S_{j+1} = -a_1 S_j - \cdots - a_j S_1 - (j+1)a_{j+1} = j - (j+1) = -1.$$

So $S_1 = \cdots = S_{i-1} = -1$. Similarly, from (3.1)

$$S_i = -a_1 S_{i-1} - \cdots - a_{i-1} S_1 - i a_i = 2i - 1.$$

Lemma 3.1. *Let $2 \leq k \leq \frac{N-1}{i} - 1$ and suppose $a_{li+n} = a_{li}$ for $1 \leq n \leq i-1$ and $0 \leq l \leq k-2$. Then we have*

$$(3.3) \quad \sum_{l=0}^{k-2} a_{li} \{ S_{(k-l)i+j+1} - S_{(k-l-1)i+j+1} \} + (ki+j+1)(a_{ki+j+1} - a_{ki+j}) \\ + \sum_{n=0}^{i-1} a_{(k-1)i+n} (S_{i+j-n+1} - S_{i+j-n}) = 0$$

for $0 \leq j \leq i-2$.

Proof. Suppose $0 \leq j \leq i-2$. From (3.1) and (3.2) we have

$$(3.4) \quad 0 = \sum_{l=0}^{k-1} \sum_{n=0}^{i-1} a_{li+n} S_{(k-l)i+j-n} + \sum_{n=0}^{j-1} a_{ki+n} S_{j-n} + (ki+j)a_{ki+j} \\ = \sum_{l=0}^{k-2} a_{li} \sum_{n=0}^{i-1} S_{(k-l)i+j-n} + \sum_{n=0}^{i-1} a_{(k-1)i+n} S_{i+j-n} \\ - \sum_{n=0}^j a_{ki+n} + (ki+j+1)a_{ki+j}.$$

Similarly,

$$(3.5) \quad 0 = \sum_{l=0}^{k-2} a_{li} \sum_{n=0}^{i-1} S_{(k-l)i+j-n+1} + \sum_{n=0}^{i-1} a_{(k-1)i+n} S_{i+j-n+1} \\ - \sum_{n=0}^j a_{ki+n} + (ki+j+1)a_{ki+j+1}.$$

Hence, on subtracting (3.5) from (3.4), we have

$$0 = \sum_{l=0}^{k-2} a_{li} \{S_{(k-l)i+j+1} - S_{(k-l-1)i+j+1}\} + (ki + j + 1)(a_{ki+j+1} - a_{ki+j}) \\ + \sum_{n=0}^{i-1} a_{(k-1)i+n}(S_{i+j-n+1} - S_{i+j-n}).$$

This proves (3.3). \square

Lemma 3.2. *Let $0 \leq k \leq \frac{N-1}{i} - 1$. Suppose $a_{li+n} = a_{li}$ for $1 \leq n \leq i-1$ and $0 \leq l \leq k$. Then*

$$(3.6) \quad S_{li+n} = -1$$

for $1 \leq n \leq i-1$ and $0 \leq l \leq k$.

Proof. We prove this by induction on k . We have proved that (3.6) is true for $k=0$. Suppose (3.6) is true for $k-1$. Then for any $0 \leq j \leq i-2$,

$$0 = a_0 \{S_{ki+j+1} - S_{(k-1)i+j+1}\} + a_{(k-1)i} \sum_{n=0}^{i-1} (S_{i+j-n+1} - S_{i+j-n}) \\ = S_{ki+j+1} + 1 + a_{(k-1)i}(S_{i+j+1} - S_{j+1}) \\ = S_{ki+j+1} + 1,$$

by (3.3). Hence $S_{ki+j+1} = -1$ for $0 \leq j \leq i-2$. \square

Theorem 3.3. *Suppose N is odd. If $a_0 = a_1 = \dots = a_{i-1} = 1$ and $a_i = -1$, then*

$$a_{li+n} = a_{li}$$

for $1 \leq n \leq i-1$ and $0 \leq l \leq \frac{N}{i} - 1$.

Proof. We first show that $S_{2k} = -1$ for $1 \leq k \leq N-1$ and hence i is odd because $S_i = 2i - 1$. Since N is odd, from Corollary 2.7,

$$P(x) = \prod_{d|N} \Phi_d^{e(d)}(x) \Phi_{2d}^{e(2d)}(x)$$

where $e(d) + e(2d) = 1$ if $d > 1$ and $e(1) = e(2) = 0$. If $1 \leq k \leq N-1$, then

$$(3.7) \quad S_{2k} = \sum_{d|N} (e(d)C_d(2k) + e(2d)C_{2d}(2k)) \\ = \sum_{d|N} (e(d) + e(2d))C_d(k) \\ = \sum_{d|N} C_d(k) - C_1(k) \\ = -1$$

where the Ramanujan sum, $C_d(k)$, is the sum of the k th powers of the primitive d th roots of unity and hence $\sum_{d|N} C_d(k)$ is the sum of k th powers of the roots of $\prod_{d|N} \Phi_d(x) = x^N - 1$ which is equal to zero when $1 \leq k \leq N-1$.

We then proceed our proof by using induction on k . Suppose

$$a_{li+n} = a_{li}$$

for $1 \leq n \leq i-1$ and $0 \leq l \leq k-1$ where $1 \leq k \leq \frac{N-1}{i} - 1$. From Lemmas 3.1 and 3.2 we have

$$(3.8) \quad S_{ki+j+1} + 1 + (ki+j+1)(a_{ki+j+1} - a_{ki+j}) = 0$$

and hence from (3.3) again

$$\begin{aligned} 0 &= a_0(S_{(k+1)i+j+1} - S_{ki+j+1}) + a_i(S_{ki+j+1} - S_{(k-1)i+j+1}) + a_{ki+j}(S_{i+1} - S_i) \\ &\quad + a_{ki+j+1}(S_i - S_{i-1}) + ((k+1)i+j+1)(a_{(k+1)i+j+1} - a_{(k+1)i+j}) \\ &= (S_{(k+1)i+j+1} - 2S_{ki+j+1} - 1) + 2i(a_{ki+j+1} - a_{ki+j}) \\ &\quad + ((k+1)i+j+1)(a_{(k+1)i+j+1} - a_{(k+1)i+j}) \\ &= S_{(k+1)i+j+1} + 1 + 2((k+1)i+j+1)(a_{ki+j+1} - a_{ki+j}) \\ (3.9) \quad &+ ((k+1)i+j+1)(a_{(k+1)i+j+1} - a_{(k+1)i+j}) \end{aligned}$$

for $0 \leq j \leq i-2$. Suppose k is even. Then in view of (3.7), $S_{ki+j+1} = -1$ if j is odd and $S_{(k+1)i+j+1} = -1$ if j is even. So from (3.8) and (3.9), we have

$$a_{ki+j+1} = a_{ki+j}$$

for $j = 1, 3, \dots, i-2$ and

$$(3.10) \quad -2(a_{ki+j+1} - a_{ki+j}) = a_{(k+1)i+j+1} - a_{(k+1)i+j}$$

for $j = 0, 2, \dots, i-3$. However, since the a_i^l s are $+1$ or -1 , (3.10) implies that

$$a_{ki+j+1} = a_{ki+j} \quad \text{and} \quad a_{(k+1)i+j+1} = a_{(k+1)i+j}$$

for $j = 0, 2, \dots, i-3$. Hence $a_{ki+n} = a_{ki}$ for $n = 1, 2, \dots, i-1$. The case k is odd can be proved in the same way. \square

Theorem 3.4. *Suppose N is odd. A Littlewood polynomial, $P(x)$, of degree $N-1$ is cyclotomic if and only if*

$$(3.11) \quad P(x) = \pm \Phi_{p_1}(\pm x) \Phi_{p_2}(\pm x^{p_1}) \cdots \Phi_{p_r}(\pm x^{p_1 p_2 \cdots p_{r-1}}),$$

where $N = p_1 p_2 \cdots p_r$ and the p_i are primes, not necessarily distinct.

Proof. It is clear that if $P(x)$ is in the form of (3.11), then $P(x)$ is a cyclotomic Littlewood polynomial. Conversely suppose $P(x)$ is a cyclotomic Littlewood polynomial. As before we may assume that $a_0 = a_1 = \cdots = a_{i-1} = 1$ and $a_i = -1$. Then we are going to prove our result by induction on N . From Theorem 3.3, we have $P(x) = P_1(x)P_2(x^i)$ where $P_1(x) = 1 + x + \cdots + x^{i-1}$ and $P_2(x)$ is a cyclotomic Littlewood polynomial of degree $< N-1$. By induction, $P_1(x)$ and $P_2(x)$ are in the form of (3.11) and hence so is $P(x)$ because the degree of $P_1(x)$ is $i-1$. This completes the proof of our theorem. \square

Corollary 3.5. *Suppose N is odd. Then $P(x)$ is cyclotomic in \mathcal{L} of degree $N-1$ if and only if*

$$P(x) = \pm \prod_{i=1}^t \frac{x^{N_i} + (-1)^{\epsilon+i}}{x^{N_{i-1}} + (-1)^{\epsilon+i}}$$

where $\epsilon = 0$ or 1 , $N_0 = 1$, $N_t = N$ and N_{i-1} is a proper divisor of N_i for $i = 1, 2, \dots, t$.

Proof. Without loss of generality, we may assume that $P(x) = 1 + x + a_2x^2 + \dots$. So from Theorem 3.4, $P(x)$ is cyclotomic in \mathcal{L} if and only if

$$\begin{aligned} P(x) &= \Phi_{p_1}(x)\Phi_{p_2}(\pm x^{p_1}) \cdots \Phi_{p_r}(\pm x^{p_1 \cdots p_{r-1}}) \\ &= \Phi_{p_1}(x) \cdots \Phi_{p_{n_1}}(x^{p_1 \cdots p_{n_1-1}})\Phi_{p_{n_1+1}}(-x^{p_1 \cdots p_{n_1}}) \cdots \Phi_{p_{n_2}}(-x^{p_1 \cdots p_{n_2-1}}) \\ (3.12) \quad &\cdots \Phi_{p_{n_{t-1}+1}}((-1)^{t-1}x^{p_1 \cdots p_{n_{t-1}}}) \cdots \Phi_{p_{n_t}}((-1)^{t-1}x^{p_1 \cdots p_{n_t-1}}) \end{aligned}$$

where $N = p_1 \cdots p_{n_t}$. Since $\Phi_p(x) = \frac{x^p - 1}{x - 1}$, (3.12) becomes

$$P(x) = \prod_{i=1}^t \frac{x^{N_i} + (-1)^i}{x^{N_{i-1}} + (-1)^i},$$

where $N_0 = 1$ and $N_i = p_1 \cdots p_{n_i}$ for $i = 1, \dots, t$. This proves our corollary. \square

Using Corollary 3.5, we can count the number of cyclotomic Littlewood polynomials of given even degree. For any positive integers N and t , define

$$r(N, t) := \#\{(N_1, N_2, \dots, N_t) : N_1 | N_2 | \cdots | N_t, 1 < N_1 < N_2 < \cdots < N_t = N\};$$

and for $i \geq 1$,

$$(3.13) \quad d_i(N) := \sum_{n|N} d_{i-1}(n)$$

where $d_0(N) = 1$.

Lemma 3.6. *For $l, t \geq 0$ and p prime, we have*

$$(3.14) \quad d_t(p^l) = \binom{l+t}{t}.$$

Proof. We prove the lemma by induction on t . Equality (3.14) is clearly true for $t = 0$ because $d_0(N) = 1$. We then suppose (3.14) is true for $t - 1$ where $t \geq 1$. Then

$$d_t(p^l) = \sum_{n|p^l} d_{t-1}(n) = \sum_{i=0}^l d_{t-1}(p^i) = \sum_{i=0}^l \binom{i+t-1}{t-1}.$$

So $d_t(p^l)$ is the coefficient of x^{t-1} in

$$\begin{aligned} &(x+1)^{t-1} + (x+1)^t + \cdots + (x+1)^{l+t-1} \\ &= (x+1)^{t-1} \left\{ \frac{(x+1)^{l+t} - 1}{x} \right\} \\ &= \frac{(x+1)^{l+t} - (x+1)^{t-1}}{x}. \end{aligned}$$

Hence $d_t(p^l)$ is the coefficient of x^t in $(x+1)^{l+t} - (x+1)^{t-1}$. Therefore, $d_t(p^l) = \binom{l+t}{t}$. \square

Since $d_t(N)$ is a multiplicative function of N , we have

Corollary 3.7. *If $N = p_1^{r_1} \cdots p_s^{r_s}$ where $r_i \geq 1$ and p_i are distinct primes, then*

$$d_t(N) = \prod_{i=1}^s \binom{r_i + t}{t}.$$

Lemma 3.8. *For any positive integers N and t , we have*

$$(3.15) \quad r(N, t) := \begin{cases} 0 & \text{if } N = 1, \\ \sum_{i=1}^t (-1)^{t-i} \binom{t}{i} d_{i-1}(N) & \text{if } N > 1. \end{cases}$$

Proof. We again prove by induction on t . It is clear from the definition that $r(1, t) = 0$ and $r(N, 1) = 1$ for any $t, N \geq 1$. We then suppose $N > 1$ and (3.15) is true for $t - 1$ where $t \geq 2$. Then

$$\begin{aligned} r(N, t) &= \sum_{\substack{N_1|N \\ N_1 > 1}} r(N/N_1, t - 1) \\ &= \sum_{\substack{N_1|N \\ N > N_1 > 1}} r(N/N_1, t - 1) \\ &= \sum_{N_1|N} \left\{ \sum_{i=1}^{t-1} (-1)^{t-i-1} \binom{t-1}{i} d_{i-1}(N/N_1) \right\} \\ &\quad - \sum_{i=1}^{t-1} (-1)^{t-i-1} \binom{t-1}{i} \{d_{i-1}(1) + d_{i-1}(N)\} \\ &= \sum_{i=2}^t (-1)^{t-i} \binom{t-1}{i-1} d_{i-1}(N) \\ &\quad - \sum_{i=1}^{t-1} (-1)^{t-i-1} \binom{t-1}{i} d_{i-1}(N) + (-1)^{t-1} \\ &= \sum_{i=1}^t (-1)^{t-i} \binom{t}{i} d_{i-1}(N) \end{aligned}$$

from (3.13) and the fact that $\binom{t-1}{i-1} + \binom{t-1}{i} = \binom{t}{i}$. \square

Corollary 3.9. *The number of cyclotomic polynomials in \mathcal{L} of degree $N - 1$ where $N = p_1^{r_1} \cdots p_s^{r_s}$, $r_i \geq 1$ and the p_i are distinct odd primes, is*

$$4 \sum_{i=1}^{r_1 + \cdots + r_s} \sum_{j=1}^i (-1)^{i-j} \binom{i}{j} \prod_{k=1}^s \binom{r_k + j - 1}{j - 1}.$$

Proof. From Corollary 3.5, the number of cyclotomic polynomials in \mathcal{L} of degree $N - 1$ is

$$4 \times \sum_{i=1}^{r_1 + \cdots + r_s} r(N, i).$$

The corollary now follows from Corollary 3.7 and Lemma 3.8. \square

4. CYCLOTOMIC LITTLEWOOD POLYNOMIALS OF ODD DEGREE.

We conjecture explicitly that Theorem 3.4 also holds for polynomials of odd degree.

Conjecture 4.1. *A Littlewood polynomial, $P(x)$, of degree $N - 1$ is cyclotomic if and only if*

$$(4.1) \quad P(x) = \pm \Phi_{p_1}(\pm x) \Phi_{p_2}(\pm x^{p_1}) \cdots \Phi_{p_r}(\pm x^{p_1 p_2 \cdots p_{r-1}}),$$

where $N = p_1 p_2 \cdots p_r$ and the p_i are primes, not necessarily distinct.

We computed up to degree 210 (except for the case $N - 1 = 191$). The computation was based on computing all cyclotomic polynomials with odd coefficients of a given degree and then checking which were actually Littlewood and seeing that this set matched the set generated by the conjecture. For example, for $N - 1 = 143$ there are 6773464 cyclotomic polynomials with odd coefficients of which 416 are Littlewood. For $N - 1 = 191$ there are 697392380 cyclotomic polynomials with odd coefficients (which was too big for our program).

We can generate all the cyclotomics with odd coefficients from Corollary 2.7 quite easily so the bulk of the work is involved in checking which ones have height 1. The set in the conjecture computes very easily recursively.

Some special cases also support the conjecture. Most notably the case where N is a power of 2. The proof is as follows. From Corollary 2.7, we have

$$P(x) = \Phi_1^{e(1)}(x) \Phi_2^{e(2)}(x) \cdots \Phi_{2^{t+1}}^{e(2^{t+1})}(x).$$

Again, we assume $a_0 = a_1 = 1$. Since $\Phi_1(x) \Phi_2(x) = x^2 - 1$ and

$$\Phi_{2^l}(x) = \Phi_2(x^{2^{l-1}})$$

for $l \geq 2$, we have $e(2) - e(1) = 1$ and hence

$$P(x) = \Phi_2(x) Q(x^2),$$

for some cyclotomic Littlewood polynomial $Q(x)$. Therefore, by induction, $P(x)$ satisfies (4.1).

ACKNOWLEDGMENT

The authors wish to thank Professor D. Boyd for his advice and support concerning this paper.

REFERENCES

- [1] P. Borwein, *Some Old Problems on Polynomials with Integer Coefficients* (To appear).
- [2] P. Borwein, T. Erdélyi & G. Kos *Littlewood-type problems on $[0, 1]$* Bull. London Math. Soc. (to appear).
- [3] P. Borwein & T. Erdélyi, *Polynomials and Polynomials Inequalities*, GTM **161**, Springer-Verlag, Berlin, 1995.
- [4] N. G. de Bruijn, *On Mahler's partition problem*, Indagationes Math., **10** (1948), 210-220.
- [5] R. Lidl & H. Niederreiter, *Finite Fields*, Addison-Wesley, Reading, 1983.
- [6] J.E. Littlewood *Some Problems in Real and Complex Analysis*, Heath Mathematical Monographs, Lexington, Mass. 1968.
- [7] K. Mahler *On two extremal properties of polynomials*, Illinois J. Math., **7** (1963), 681-701.
- [8] T. W. Hungerford, *Algebra*, GTM **73**, Springer-Verlag, Berlin, 1980.

DEPARTMENT OF MATHEMATICS AND STATISTICS, SIMON FRASER UNIVERSITY, BURNABY, B.C., CANADA V5A 1S6

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, B.C., CANADA V6T 1Z2 AND DEPARTMENT OF MATHEMATICS AND STATISTICS, SIMON FRASER UNIVERSITY, BURNABY, B.C., CANADA V5A 1S6