

**LOWER BOUNDS FOR THE MERIT  
FACTORS OF TRIGONOMETRIC  
POLYNOMIALS FROM LITTLEWOOD CLASSES**

PETER BORWEIN AND TAMÁS ERDÉLYI

ABSTRACT. With the notation  $K := \mathbb{R} \pmod{2\pi}$ ,

$$\|p\|_{L_\lambda(K)} := \left( \int_K |p(t)|^\lambda dt \right)^{1/\lambda} \quad \text{and} \quad M_\lambda(p) := \left( \frac{1}{2\pi} \int_K |p(t)|^\lambda dt \right)^{1/\lambda}$$

we prove the following result.

**Theorem 1.** *Assume that  $p$  is a trigonometric polynomial of degree at most  $n$  with real coefficients that satisfies*

$$\|p\|_{L_2(K)} \leq An^{1/2} \quad \text{and} \quad \|p'\|_{L_2(K)} \geq Bn^{3/2}.$$

Then

$$M_4(p) - M_2(p) \geq \varepsilon M_2(p)$$

with

$$\varepsilon := \left( \frac{1}{111} \right) \left( \frac{B}{A} \right)^{12}.$$

We also prove that

$$M_\infty(1+2p) - M_2(1+2p) \geq (\sqrt{4/3} - 1)M_2(1+2p)$$

and

$$M_2(p) - M_1(p) \geq 10^{-31} M_2(p)$$

for every  $p \in \mathcal{A}_n$ , where  $\mathcal{A}_n$  denotes the collection of all trigonometric polynomials of the form

$$p(t) := p_n(t) := \sum_{j=1}^n a_j \cos(jt + \alpha_j), \quad a_j = \pm 1, \quad \alpha_j \in \mathbb{R}.$$

---

1991 *Mathematics Subject Classification.* Primary: 41A17.

*Key words and phrases.* trigonometric polynomials, merit factor, unimodular trigonometric polynomials, Littlewood class..

Research of P. Borwein is supported by MITACS and by NSERC of Canada. Research of T. Erdélyi is supported, in part, by NSF under Grant No. DMS-0070826.

## INTRODUCTION

We give shorter and more direct proofs of some of the main results from Littlewood's papers [Li-61], [Li-62], [Li-66a], [Li-66b], and [Li-68]. There are two reasons for doing this. First our approaches are, we believe, much easier, and secondly they lead to explicit constants. Littlewood himself remarks that his methods were "extremely indirect." Motivation and discussion of these types of results may be found in [Bo-02]. Kahane's paper [Ka-85] is also central among those related to the subject of this paper.

## 2. NEW RESULTS

We use the notation  $K := \mathbb{R} \pmod{2\pi}$ . Let

$$\|p\|_{L_\lambda(K)} := \left( \int_K |p(t)|^\lambda dt \right)^{1/\lambda} \quad \text{and} \quad M_\lambda(p) := \left( \frac{1}{2\pi} \int_K |p(t)|^\lambda dt \right)^{1/\lambda}.$$

**Theorem 1.** *Assume that  $p$  is a trigonometric polynomial of degree at most  $n$  with real coefficients that satisfies*

$$(1) \quad \|p\|_{L_2(K)} \leq An^{1/2}$$

and

$$(2) \quad \|p'\|_{L_2(K)} \geq Bn^{3/2}.$$

Then

$$M_4(p) - M_2(p) \geq \varepsilon M_2(p)$$

with

$$\varepsilon := \left( \frac{1}{111} \right) \left( \frac{B}{A} \right)^{12}.$$

Let the Littlewood class  $\mathcal{A}_n$  be the collection of all trigonometric polynomials of the form

$$p(t) := p_n(t) := \sum_{j=1}^n a_j \cos(jt + \alpha_j), \quad a_j = \pm 1, \quad \alpha_j \in \mathbb{R}.$$

Note that for the Littlewood class  $\mathcal{A}_n$  we have

$$\left( \frac{B}{A} \right)^{12} = 3^{-6}.$$

**Corollary 2.** *We have*

$$M_4(p) - M_2(p) \geq \frac{M_2(p)}{80920}$$

for every  $p \in \mathcal{A}_n$ . The merit factor

$$\left( \frac{M_4^A(p)}{M_2^A(p)} - 1 \right)^{-1}$$

is bounded above by 20230 for every  $p \in \mathcal{A}_n$ .

If  $Q_n$  is a polynomial of degree  $n$  of the form

$$Q_n(z) = \sum_{k=0}^n a_k z^k, \quad a_k \in \mathbb{C},$$

and the coefficients  $a_k$  of  $Q_n$  satisfy

$$a_k = \bar{a}_{n-k}, \quad k = 0, 1, \dots, n,$$

then we call  $Q_n$  a *conjugate-reciprocal* polynomial of degree  $n$ . We say that the polynomial  $Q_n$  is unimodular, if  $|a_k| = 1$  for each  $k = 0, 1, 2, \dots, n$ . Note that if  $p \in \mathcal{A}_n$ , then

$$1 + 2p(t) = e^{int} Q_{2n}(e^{it})$$

with a conjugate-reciprocal unimodular polynomial  $Q_{2n}$  of degree exactly  $2n$ . One can ask how flat a conjugate reciprocal unimodular polynomial can be. Here we reprove a result of Erdős [Er-62]. His proof is much longer and his constant  $\varepsilon > 0$  is unspecified. This result has already been recorded in [Er-01].

**Theorem 3.** *Let  $\partial D$  denote the unit circle. Let  $P$  be a conjugate reciprocal unimodular polynomial of degree  $n$ . Then*

$$\max_{z \in \partial D} |P(z)| \geq (1 + \varepsilon) \sqrt{n + 1}$$

with  $\varepsilon := \sqrt{4/3} - 1$ . As a consequence, we have

$$M_\infty(1 + 2p) - M_2(1 + 2p) \geq (\sqrt{4/3} - 1) M_2(1 + 2p)$$

for every  $p \in \mathcal{A}_n$ .

In our next theorem we give the numerical value of an unspecified constant appearing in another main result of Littlewood. In the proof we will need to refer to only a two-page-long (very clever) piece of Littlewood's paper [Li-66a].

**Theorem 4.** *We have*

$$M_2(p) - M_1(p) \geq 10^{-31} M_2(p)$$

for every  $p \in \mathcal{A}_n$ .

Based on the fact that for a fixed trigonometric polynomial  $p$  the function

$$\lambda \rightarrow \lambda \log(M_\lambda(p))$$

is a convex function on  $[0, \infty)$ , we can state explicit numerical values of certain unspecified constants in some other related Littlewood results. For example, as a consequence of Theorem 4, we have

**Theorem 5.** *We have*

$$\log(M_\lambda(p)) - \log(M_2(p)) \geq \frac{\lambda - 2}{\lambda} \log\left(\frac{1}{1 - 10^{-31}}\right), \quad \lambda > 2,$$

and

$$\log(M_2(p)) - \log(M_\lambda(p)) \geq \frac{2 - \lambda}{\lambda} \log\left(\frac{1}{1 - 10^{-31}}\right), \quad 1 \leq \lambda < 2,$$

for every  $p \in \mathcal{A}_n$ .

### 3. PROOFS

*Proof of Theorem 1.* For the sake of brevity let  $\mu_n := \mu_n(p) = M_2(p)$ . Note that Bernstein's inequality in  $L_2(K)$  implies  $B \leq A$ . Without loss of generality we may assume that

$$(3) \quad \|p\|_{L_4(K)}^4 \leq 2\pi \frac{33}{32} \mu_n^4.$$

Then by the Bernstein Inequality for trigonometric polynomials in  $L_4(K)$  we can deduce that

$$\|p'\|_{L_4(K)} \leq n \|p\|_{L_4(K)} \leq n \left(\frac{33}{32}\right)^{1/4} (2\pi)^{1/4} \mu_n \leq \pi^{-1/4} \left(\frac{33}{64}\right)^{1/4} A n^{3/2}.$$

Hence, combining this with (2) and Hölder's Inequality, we obtain

$$B^2 n^3 \leq \int_K |p'(t)|^2 dt \leq \|p'\|_{L_1(K)}^{2/3} \|p'\|_{L_4(K)}^{4/3} \leq \|p'\|_{L_1(K)}^{2/3} \pi^{-1/3} \left(\frac{33}{64}\right)^{1/3} A^{4/3} n^2.$$

Therefore

$$\frac{\pi^{1/3} \left(\frac{64}{33}\right)^{1/3} B^2}{A^{4/3}} n \leq \|p'\|_{L_1(K)}^{2/3},$$

that is

$$\frac{\pi^{1/2} \left(\frac{64}{33}\right)^{1/2} B^3}{A^2} n^{3/2} \leq \|p'\|_{L_1(K)}.$$

Combining this with (1), we have

$$\gamma n \mu_n \leq \gamma n \frac{A}{(2\pi)^{1/2}} \sqrt{n} \leq \frac{\pi^{1/2} \left(\frac{64}{33}\right)^{1/2} B^3}{A^2} n^{3/2} \leq \|p'\|_{L_1(K)}$$

with

$$\gamma := \frac{\left(\frac{128}{33}\right)^{1/2} \pi B^3}{A^3}.$$

Now let

$$E := E(n, p, \gamma) := \left\{ t \in [0, 2\pi) : (|p(t)| - \mu_n)^2 \geq \left(\frac{\gamma \mu_n}{16}\right)^2 \right\}.$$

Then using Hölder's Inequality and then Bernstein's Inequality for trigonometric polynomials in  $L_2(K)$ , we can deduce that

$$\begin{aligned} \gamma n \mu_n &\leq \int_0^{2\pi} |p'(t)| dt \leq \int_{[0, 2\pi] \setminus E} |p'(t)| dt + \int_E |p'(t)| dt \\ &\leq 2 \cdot (2n) \cdot \frac{2\gamma}{16} \mu_n + \int_E |p'(t)| dt \leq \frac{\gamma}{2} n \mu_n + \sqrt{m(E)} \left( \int_E |p'(t)|^2 dt \right)^{1/2} \\ &\leq \frac{\gamma}{2} n \mu_n + \sqrt{m(E)} n \left( \int_0^{2\pi} |p(t)|^2 dt \right)^{1/2} \leq \frac{\gamma}{2} n \mu_n + \sqrt{m(E)} n (2\pi)^{1/2} \mu_n. \end{aligned}$$

Hence

$$\frac{\gamma}{2} n \mu_n \leq \sqrt{m(E)} n (2\pi)^{1/2} \mu_n,$$

that is

$$(4) \quad \beta := \frac{\gamma^2}{8\pi} \leq m(E).$$

So we have

$$\begin{aligned} 2\pi(M_4(p)^4 - M_2(p)^4) &= \|p\|_{L_4(K)}^4 - 2\pi\mu_n^4 \\ &= \int_0^{2\pi} (p(t)^2 - \mu_n^2)^2 dt \\ &\geq m(E) \left( \frac{\gamma\mu_n}{16} \right)^2 \mu_n^2 \geq \frac{\gamma^2}{8\pi} \left( \frac{\gamma\mu_n}{16} \right)^2 \mu_n^2 \\ &= \frac{1}{2^{11}\pi} \left( \frac{128}{33} \right)^2 \pi^4 \left( \frac{B}{A} \right)^{12} \mu_n^4. \end{aligned}$$

Combining this with (3) we obtain

$$M_4(p) - M_2(p) \geq 2^{-14} \left( \frac{B}{A} \right)^{12} \left( \frac{128}{33} \right)^2 \pi^2 M_2(p),$$

and the theorem is proved Here we used

$$M_4(p)^4 - M_2(p)^4 \geq (M_4(p) - M_2(p))4M_2(p)^3,$$

which is a consequence of the Mean Value Theorem. Note that

$$\frac{1}{111} \leq 2^{-14} \left( \frac{128}{33} \right)^2 \pi^2 \leq \frac{1}{110}.$$

□

*Proof of Theorem 3.* Let  $P$  be a conjugate reciprocal unimodular polynomial of degree  $n$ . To prove the statement, observe that Malik's inequality [MMR, p. 676] gives

$$\max_{z \in \partial D} |P'(z)| \leq \frac{n}{2} \max_{z \in \partial D} |P(z)|.$$

(Note that the fact that  $P$  is conjugate reciprocal improves the Bernstein factor for  $P$  on  $\partial D$  from  $n$  to  $n/2$ .) Using the fact that each coefficient of  $P$  is of modulus 1, then applying Parseval's formula and Malik's inequality, we obtain

$$2\pi \frac{n^2(n+1)}{3} \leq 2\pi \frac{n(n+1)(2n+1)}{6} = \int_{\partial D} |P'(z)|^2 |dz| \leq 2\pi \left(\frac{n}{2}\right)^2 \max_{z \in \partial D} |P(z)|^2,$$

and

$$\max_{z \in \partial D} |P(z)| \geq \sqrt{4/3} \sqrt{n+1}$$

follows.  $\square$

*Proof of Theorem 4.* Let  $p \in \mathcal{A}_n$ . For the sake of brevity let  $\mu_n := \mu_n(p) = M_2(p)$ . Let  $N(p, v)$  be the number of real roots of  $p - v\mu_n = 0$  in  $(-\pi, \pi)$ . Littlewood proves (see Theorem 1 (i) of [Li-66a]) that if  $p \in \mathcal{A}_n$  and

$$\frac{1}{2\pi} \int_0^{2\pi} |p(t)| dt = c\mu_n,$$

then

$$N(p, v) \geq 2^{-16} c^{11} n, \quad |v| \leq 2^{-5} c^3.$$

The reader may wish to find this lower bound hidden in the proof of Theorem 1 (i) of Littlewood's paper [Li-66a]. Hence, by estimating the total variation in the usual way,  $\gamma n \mu_n \leq \|p'\|_{L_1(K)}$  with  $\gamma := 2^{-20} c^{14}$ . If  $c \leq 2^{-1/14}$ , then the proof of the theorem is finished. If  $c \geq 2^{-1/14}$ , then  $\gamma \geq 2^{-21}$ , so in the sequel we may assume that  $\gamma \geq 2^{-21}$  holds. Now let

$$E := E(n, p, \gamma) := \left\{ t \in [0, 2\pi) : (|p(t)| - \mu_n)^2 \geq \left(\frac{\gamma \mu_n}{16}\right)^2 \right\}.$$

Estimating the total variation of  $p$  on  $[0, 2\pi] \setminus E$ , then using Hölder's Inequality and then Bernstein's Inequality for trigonometric polynomials in  $L_2(K)$ , we can deduce that

$$\begin{aligned} \gamma n \mu_n &\leq \int_0^{2\pi} |p'(t)| dt \leq \int_{[0, 2\pi] \setminus E} |p'(t)| dt + \int_E |p'(t)| dt \\ &\leq 2 \cdot (2n) \cdot \frac{2\gamma}{16} \mu_n + \int_E |p'(t)| dt \leq \frac{\gamma}{2} n \mu_n + \sqrt{m(E)} \left( \int_E |p'(t)|^2 dt \right)^{1/2} \\ &\leq \frac{\gamma}{2} n \mu_n + \sqrt{m(E)} n \left( \int_0^{2\pi} |p(t)|^2 dt \right)^{1/2} \leq \frac{\gamma}{2} n \mu_n + \sqrt{m(E)} n (2\pi)^{1/2} \mu_n. \end{aligned}$$

Hence

$$\frac{\gamma}{2} n \mu_n \leq \sqrt{m(E)} n (2\pi)^{1/2} \mu_n,$$

that is

$$\beta := \frac{\gamma^2}{8\pi} \leq m(E).$$

So we have

$$\begin{aligned} 4\pi((M_2(p))^2 - M_2(p)M_1(p)) &= \int_0^{2\pi} (|p(t)| - \mu_n)^2 dt \\ &\geq m(E) \left(\frac{\gamma\mu_n}{16}\right)^2 \geq \frac{\gamma^2}{8\pi} \left(\frac{\gamma\mu_n}{16}\right)^2 \\ &= \frac{\gamma^4}{2^{11}\pi} \mu_n^2 \geq 2^{-95} \pi^{-1} \mu_n^2. \end{aligned}$$

This implies

$$M_2(p) - M_1(p) \geq 2^{-97} \pi^{-2} M_2(p),$$

and the theorem is proved.  $\square$

#### ACKNOWLEDGEMENTS

The authors wish to thank the referee, Szilárd Révész, for his careful reading of the original version of this manuscript. The constant  $1/111$  in Theorem 1 would not be the same without him.

#### REFERENCES

- [Bo-02] P. Borwein, *Computational Excursions in Analysis and Number Theory*, Springer-Verlag, New York, 2002.
- [Er-01] T. Erdélyi The phase problem of ultraflat unimodular polynomials: the resolution of the conjecture of Saffari, *Math. Annalen* **321** (2001), 905–924.
- [Er-62] P. Erdős, *An inequality for the maximum of trigonometric polynomials*, *Annales Polonica Math.* **12** (1962), 151–154.
- [Ka-85] J-P. Kahane, *Sur les polynômes à coefficients unimodulaires*, *Bull. London Math. Soc* **12** (1980), 321–342.
- [Li-61] J.E. Littlewood, *On the mean values of certain trigonometrical polynomials*, *J. London Math. Soc.* **36** (1961), 307–334.
- [Li-64] J.E. Littlewood, *On the real roots of real trigonometrical polynomials (II)*, *J. London Math. Soc.* **39** (1964), 511–552.
- [Li-66a] J.E. Littlewood, *The real zeros and value distributions of real trigonometrical polynomials*, *J. London Math. Soc.* **41** (1966), 336–342.
- [Li-66b] J.E. Littlewood, *On polynomials  $\sum \pm z^m$  and  $\sum e^{\alpha_m i} z^m$ ,  $z = e^{\theta i}$* , *J. London Math. Soc.* **41** (1966), 367–376.
- [Li-68] J.E. Littlewood, *Some Problems in Real and Complex Analysis*, Heath Mathematical Monographs, Lexington, Massachusetts, 1968.
- [MMR] Milovanović, G.V., D.S. Mitrinović, & Th.M. Rassias, *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*, World Scientific, Singapore, 1994.

DEPARTMENT OF MATHEMATICS AND STATISTICS, SIMON FRASER UNIVERSITY, BURNABY, B.C., CANADA V5A 1S6 (P. BORWEIN)

*E-mail address:* pborwein@cecm.sfu.ca (Peter Borwein)

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS  
77843 (T. ERDÉLYI)

*E-mail address:* terdelyi@math.tamu.edu (Tamás Erdélyi)